



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Antimicrobial host defence peptides: functions and clinical potential

### Citation for published version:

Mookherjee, N, Anderson, MA, Haagsman, HP & Davidson, DJ 2020, 'Antimicrobial host defence peptides: functions and clinical potential', *Nature Reviews Drug Discovery*. <https://doi.org/10.1038/s41573-019-0058-8>

### Digital Object Identifier (DOI):

[10.1038/s41573-019-0058-8](https://doi.org/10.1038/s41573-019-0058-8)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Nature Reviews Drug Discovery

### Publisher Rights Statement:

Author's peer reviewed manuscript as accepted for publication.

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Antimicrobial host defence peptides: Functions and clinical potential

Neeloffer Mookherjee<sup>1,\*</sup>, Marilyn A. Anderson<sup>2</sup>, Henk P. Haagsman<sup>3</sup> and Donald J. Davidson<sup>4</sup>

## Affiliations

<sup>1</sup>Manitoba Centre for Proteomics and Systems Biology, Departments of Internal Medicine and Immunology, University of Manitoba, 715 McDermot Ave, Winnipeg, MB, R3X1H8, Canada.

<sup>2</sup>La Trobe Institute for Molecular Science, La Trobe University, Melbourne, Vic. 3086, Australia.

<sup>3</sup>Division Molecular Host Defence, Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 1, 3584 CL, Utrecht, The Netherlands

<sup>4</sup>University of Edinburgh Centre for Inflammation Research, Queen's Medical Research Institute, Edinburgh BioQuarter, 47 Little France Crescent, Edinburgh EH16 4TJ, Scotland, UK

## \*Corresponding author:

Neeloffer Mookherjee

neeloffer.mookherjee@umanitoba.ca

**Keywords:** Antimicrobial peptides, Cathelicidins, Defensins, Inflammation, Immunotherapy.

## Abstract

Cationic host defence peptides (CHDP), also known as antimicrobial peptides, are naturally-occurring peptides which can combat infections through their direct microbicidal properties and/or by influencing the host's immune responses. The unique ability of CHDP to control infections as well as resolve harmful inflammation has generated interest in harnessing the properties of these peptides to develop new therapies for infectious diseases, chronic inflammatory disorders and wound healing. Various strategies have been employed to design synthetic optimized peptides, with negligible toxicity. Here, we focus on the progress made in understanding the scope of functions of CHDP and the emerging potential clinical applications of CHDP-based therapies.

## 1 Introduction

2 The global increase in multi-drug resistant pathogens along with a steady decline in the discovery of  
3 new antibiotics, underscores the need for new therapies to control infections. In addition, common  
4 treatments used for many chronic inflammatory diseases such as corticosteroids can increase  
5 susceptibility to infections<sup>1,2</sup>, including antibiotic recalcitrant pathogens<sup>3</sup>. There is therefore an urgent  
6 need for alternative strategies that can both kill pathogens as well as resolve harmful inflammation.  
7 One promising approach has emerged from the identification of Cationic Host Defence Peptides  
8 (CHDP), which can control infections either by their direct microbicidal properties and/or by  
9 modulating host immune responses, while also exhibiting the capacity to limit enhanced inflammation.

10 CHDP, also known as antimicrobial peptides, were first described in the 1960s by Kiss and Michl in  
11 the speckled frog<sup>4</sup>. Several seminal studies in the 1980s further defined antimicrobial peptides, notably  
12 cecropin isolated from moths by the Boman group<sup>5</sup>, defensins from human neutrophils by Ganz and  
13 Lehrer<sup>6</sup>, and magainins from amphibians by Zasloff et al<sup>7</sup>. CHDP are now known to be expressed  
14 across a diverse range of species, from microbes, plants, invertebrates to more complex amphibians and  
15 mammals.

16 CHDP are typically amphipathic small peptides with less than 50 amino acids and a net positive  
17 charge of +2 to +9 at physiological pH. However, these natural peptides differ significantly in sequence  
18 and structure, and are broadly classified into four structural groups based on conformation;  $\alpha$ -helical  
19 linear peptides, those with  $\beta$ -sheet with disulphide bridges, cyclic peptides and peptides with extended  
20 flexible loop structures.

21 Initial research in the field was focused on the antimicrobial functions of this family of peptides,  
22 with a drive to discover new antibiotic-like therapies based on the naturally-occurring peptides.  
23 However, studies published in the 2000s established that the microbicidal action of some of these  
24 peptides e.g. the human cathelicidin LL-37 and  $\beta$ -defensins, were severely impaired under  
25 physiological salt concentrations (particularly by divalent cations) and in the presence of host factors  
26 such as anionic polysaccharides, apolipoproteins, DNA, f-actin and glycosaminoglycans<sup>8,9</sup>. The  
27 relatively poor direct microbicidal capabilities of these peptides under physiological conditions, at the  
28 low, naturally occurring concentrations observed *in vivo*, did not adequately explain their contribution  
29 in host defense against infections<sup>10,11</sup>. Subsequently, research over the last two decades has  
30 demonstrated that CHDP exhibit a wide range of functions beyond microbicidal activity, including the  
31 ability to influence immune responses, which contribute to their antimicrobial impact (reviewed in<sup>12-15</sup>).

This has led to the adoption of the current name for this family of peptides, Cationic Host Defence Peptides, which encompasses the wide range of described functions.

Over the last three decades there has been substantial interest in therapeutically harnessing CHDP, with more than 5000 papers published in this area of research since 2017 alone. These include the examination of potential clinical uses for CHDP ranging from infections including multidrug-resistant bacteria<sup>16-19</sup>, to chronic inflammatory diseases such as arthritis<sup>20</sup>, asthma<sup>21</sup> and colitis<sup>22</sup>, as well as some cancers<sup>23</sup>. Peptide-based therapeutics currently in clinical trials are primarily for the treatment of infections such as respiratory, oral and catheter-related infections, and for wound healing (see <http://dramp.cpu-bioinfor.org/browse/ClinicalTrialsData.php>).

This review will provide an overview of current understanding of the scope of functions of CHDP, primarily from eukaryotes. Emerging therapeutic applications of these peptides, current clinical trials and the associated clinical developmental challenges will be discussed. Although there is increasing interest in the development of non-peptide mimics of CHDP for therapeutic application, such as peptoid analogs (reviewed in<sup>24</sup>), a comprehensive discussion of these approaches is beyond the scope of this review.

## **[H1] Naturally occurring CHDP**

The antimicrobial peptide database has catalogued more than 2600 natural antimicrobial peptides, including those annotated as immunomodulatory<sup>25</sup>. The major families of CHDP from eukaryotes that are of interest from a drug discovery perspective are summarized below.

## **[H2] Vertebrate CHDP**

CHDP from vertebrates have an essential role in the first line of defense against microbial pathogens. Upon infection, CHDP can kill pathogens through diverse mechanisms<sup>26-31</sup> (discussed below), acting rapidly and directly on the pathogen when present in high local concentrations, or indirectly to modify components of host defense. These peptides exhibit immunomodulatory activities that can be either pro- or anti-inflammatory depending on the phase of the infection (see below)<sup>12-14,29</sup>. CHDP from vertebrates are amphipathic peptides containing amino acids with hydrophilic and hydrophobic side chains at opposite sides of the molecules. These CHDP can interact with the negatively charged membranes of bacteria. The two main classes of CHDP in vertebrates, the defensins and cathelicidins (Fig. 1a), are produced as prepropeptides which are cleaved to yield mature active peptides.

Defensins have a common  $\beta$ -sheet core stabilized with three disulphide bridges between 6 conserved cysteine residues, and are subdivided into  $\alpha$ -,  $\beta$ - and  $\theta$ -defensins based on the linkage of cysteine residues. The genes encoding  $\alpha$ - and  $\beta$ -defensins are adjacent on chromosome 8p23.1 and probably have common ancestry, with  $\alpha$ -defensins thought to have evolved by gene duplication, and only found in some mammals, primarily primates and rodents<sup>32</sup>. Several of the human  $\alpha$ -defensins are highly expressed in neutrophils, and other  $\alpha$ -defensins are produced and secreted by Paneth cells in the small intestine<sup>6,33,34</sup>. The  $\beta$ -defensins are ubiquitous and present in all vertebrates. The human genome has >30 genes coding for  $\beta$ -defensins, and mice have even more genes coding for these peptides. These peptides are mainly produced in epithelial cells<sup>33,35</sup>. Finally, the cyclic  $\theta$ -defensins arose from  $\alpha$ -defensins after the divergence of primates and have been purified from the leukocytes of rhesus macaques and baboons. These molecules are the only cyclic peptides found in mammals and exhibit antiviral activity. Thus varying between different species, evidence exists for sequence divergence by both positive and negative selection of mammalian defensin genes following ancestral gene duplication<sup>36-38</sup>.

Cathelicidins are produced as prepropeptides containing an N-terminal signal peptide, a cathelin-like domain, and the C-terminal mature peptide. The pro-cathelin-like domain is cleaved off primarily by serine proteases once the peptide is secreted<sup>39,40</sup>. These peptides are named after the conserved cathelin-like domain in the pro-peptide which is the common denominator of this family of peptides, as the active mature cathelicidins do not share sequence similarities. Most cathelicidins are  $\alpha$ -helical (23-37 amino acids), amphipathic, cationic peptides with a hydrophobic surface enabling interaction and perturbation of membranes with anionic surfaces. The only human cathelicidin LL-37 is the most well studied peptide in this family. LL-37 is an  $\alpha$ -helical peptide and is one of several cleavage products of hCAP18, the product of the only human cathelicidin gene *CAMP*<sup>26</sup>. Other  $\alpha$ -helical cathelicidins include the sole mouse cathelicidin CRAMP<sup>41</sup> and chicken CATH-2<sup>29</sup>. Another class of cathelicidins comprises  $\beta$ -hairpin peptides (12-18 residues) such as the bovine bactenecin that have one intramolecular disulphide bond. Cathelicidins such as protegrins (16-18 residues) have two intramolecular disulphide bonds and adopt  $\beta$ -sheet structures. Finally, linear cathelicidins (13-39 residues) are enriched in specific amino acids e.g. tryptophan-enriched bovine indolicidin and proline-rich porcine PR-39. Cathelicidins are immunomodulatory antimicrobials with an important role in the regulation of the inflammatory response<sup>29-31</sup>.

Several other vertebrate CHDP are also being explored as potential alternatives to antibiotics. Among these are histatins, which are histidine-rich CHDP constituents of saliva of mammals. Histatins

are less amphipathic than cathelicidins or defensins and may have different modes of action compared to other vertebrate CHDP as illustrated by their antifungal activity<sup>42</sup>. Finally, several CHDP produced by amphibian skin granular glands, such as magainin-2, have been used as prototypes for the development of novel antimicrobials<sup>43</sup>.

## **[H2] Invertebrate, plant and fungal CHDP**

CHDP are a vital component of the innate immune system of all eukaryotic organisms, including plants, fungi and invertebrates which lack an adaptive immune system. These CHDP, similar to those from vertebrates, are also small amphipathic peptides with an overall positive net charge<sup>44,45</sup>. These peptides fall into several diverse and distinct structural groups, including  $\alpha$ -helical peptides,  $\beta$ -sheet peptides, peptides with mixed  $\alpha$ -helical and  $\beta$ -sheet structures, extended peptides and peptides enriched in specific amino acids (Fig. 1b). Their classification is generally dependent on the number, spacing and connectivity of cysteine residues as well as structure which is normally highly conserved within a CHDP family.

In plants, gene duplication and rapid evolution has produced large families of antimicrobial peptides within the same organism<sup>46</sup>. Genome sequencing has revealed that plants such as *Arabidopsis* and *Medicago* have up to 300 defensin or defensin-like sequences. Apart from the cysteine residues and a glycine residue which are required for the defensin fold, these are highly diverse wherein the sequence diversity is displayed on the surface loops presented on conserved scaffolds. This may explain the diversity of biological functions defined for this family of proteins<sup>47</sup>. Plant defensins are generally potent antifungal molecules. However, plant defensins with antibacterial and insecticidal activity have also been described, along with those with roles in cell signalling, self- non-self-recognition during sexual reproduction, ion channel perturbation and enzyme inhibition<sup>46,47</sup>. Similar to the mammalian defensins, these also exhibit other functions, including roles in innate immunity, in addition to antimicrobial activity<sup>33,48</sup>. Although once considered an example of evolution of an ancient innate immunity molecule, it is now evident that mammalian and plant defensins evolved independently and are an excellent example of convergent evolution<sup>46</sup>.

## **[H1] Synthetic peptides derived from CHDP**

The vast repertoire of sequences and structures of natural antimicrobial peptides allows for the design and development of novel therapeutic peptide analogues or peptidomimetics. Various approaches are being applied in the design of new drug candidates based on CHDP; such as single amino acid

substitution, using internal segments of CHDP to derive smaller peptides, as well as *in silico* methods to predict new synthetic peptides based on the understanding of structure-function relationships<sup>49-51</sup>.

## **[H2] Synthetic innate defence regulator (IDR) peptides**

Early approaches to the development of CHDP as therapeutics focused heavily on optimizing the direct microbicidal properties of these peptides. However, it has become clear that strategies adopted to enhance *in vitro* microbicidal functions often resulted in peptides with increased levels of cytotoxicity. In addition, naturally occurring CHDP can also exhibit other functions undesirable for drug development such as an ability to induce mast cell degranulation, with release of histamine and prostaglandin, as well as activation of complement factors<sup>52,53</sup>. Therefore, in the last decade, effective strategies have focused on designing synthetic peptides from sequences of CHDP to optimize antimicrobial functions *in vivo*, through a combination of some microbicidal activity along with beneficial immunomodulatory functions, in the absence of cytotoxic effects.

Innate defence regulator (IDR) peptides are such small synthetic cationic peptides derived from natural CHDP, which have been screened and optimized for immunomodulatory functions<sup>12,54</sup>. Libraries of IDR peptides have been generated by screening overlapping fragments representing internal sequences of CHDP, and by single amino acid substitutions<sup>49-51</sup>. Most of the IDR peptides published to date are derivatives of the bovine CHDP batenecin<sup>21,49,54</sup>. IDR peptides are non-immunogenic and do not have the potentially adverse effects associated with some natural CHDP. In general, IDR peptides have a modest direct effect on the pathogen, yet can control infection *in vivo* and reduce inflammation, as shown in a variety of infection models<sup>54-56</sup>. The beneficial use of IDR peptides for the control of infections is now well established in preclinical models of a wide range of pathogens, including multidrug resistant bacteria, viruses, parasites, and antibiotic-recalcitrant bacterial biofilms<sup>54,56-60</sup>. IDR peptides also exhibit adjuvant functions to enhance mucosal immunity and antigen-specific humoral response<sup>61-64</sup>, and are effective in controlling inflammation and related physiological outcomes in models of sterile inflammation<sup>65</sup> and chronic inflammatory diseases<sup>21</sup>. A distinct advantage of IDR peptide-based therapy, compared to current anti-inflammatory therapeutics, is the potential to resolve infections along with the ability to control inflammation<sup>54,57</sup>.

## **[H2] Cryptic and synthetic peptides**

Several mammalian cationic proteins like histones, lactoferrin, thrombin and their cleavage products have antimicrobial activities. Histones and fragments thereof have a wide range of antimicrobial

activities<sup>66</sup>. These are not restricted to nucleosomes but are also found in the cytoplasm of cells and can be released upon activation. Indeed, up to 70% of the proteins of neutrophil extracellular traps (NETs) comprise histones and histone fragments<sup>67</sup>. Lactoferrin is another example of a protein that yields antimicrobial peptides upon proteolytic degradation *in vivo*. This protein is not confined to milk, it is produced by several tissues and has several biological activities related to host defense, including antimicrobial activity. Its proteolysis in the gastrointestinal tract produces fragments that are more active than the native protein<sup>68</sup>. Proteolytic digestion with pepsin releases lactoferricin, a peptide derived from the N-terminal region of lactoferrin. Other active fragments described are Lf(1-11) and lactoferrampin<sup>69</sup>. Similarly, synthetic molecules based on human  $\theta$ -defensin pseudogenes, termed retrocyclins, have been developed as potential therapeutics for example as antivirals<sup>70</sup>. These observations that antimicrobial and/or immunomodulatory peptides can be released from larger proteins during inflammation or infection has led to the exciting concept of developing synthetic CHDP as prodrugs in which the N-terminus of the peptide is modified by a linker and a negatively charged promoiety. Such prodrugs can be selectively cleaved by a disease-associated enzyme, thereby targeting peptide activity to pathologically affected parts of the body<sup>71</sup>. In addition, these discoveries have led to bioinformatics approaches to identify segments of secreted proteins that could be developed as novel antimicrobial peptides of human origin. Bioinformatic tools have been developed to find those cryptic sequences and the peptides thus defined are known as cryptic peptides<sup>72</sup>. Among the identified and tested cryptic peptides are those from two apolipoproteins<sup>73</sup>. Other researchers have developed entirely novel antimicrobial peptides by high throughput screening of libraries of peptides that have been synthesized semi-randomly<sup>74,75</sup> or by rational design<sup>51</sup>.

## **[H1] Antimicrobial actions of CHDP**

The antimicrobial effects of CHDP on a wide range of pathogens have provided the impetus for the development of these peptides as broad-spectrum antimicrobials. Related mechanisms of action are diverse and appear to be dependent on the microbial pathogen.

## ***[H2] Antibacterial activity***

The bacterial membrane is the main target for most cationic peptides. Bacterial membranes are negatively charged because of the presence of anionic lipids, lipopolysaccharides (LPS; in Gram-negative bacteria) or teichoic acids (in Gram-positive bacteria). These negatively charged molecules initiate an electrostatic interaction with cationic compounds which explains the preference of CHDP for



1 bacterial membranes compared to the membranes of cells from plants, invertebrates and vertebrates.  
2 The amphipathic nature of CHDP is important for the membrane-destabilizing properties of these  
3 peptides. CHDP can bind to the bacterial inner membrane leading to penetration of the peptide, leakage  
4 of bacterial cell contents and cell death<sup>27,28</sup> (Fig. 2). In Gram-negative bacteria, interaction of CHDP  
5 with LPS results in perturbation of the outer membrane and this has been defined as a primary mode of  
6 action for the antimicrobial activity of CHDP<sup>76</sup>. Four main models of (inner) membrane perturbation  
7 have been proposed: aggregate, toroidal, barrel-stave and carpet model<sup>76</sup> (Fig. 2). It should be  
8 cautioned that these proposed mechanisms of action stem largely from experiments with model  
9 membranes. As the interaction of CHDP with bacterial membranes does not involve a specific target, it  
10 has been speculated that the development of microbial resistance is unlikely. Indeed, CHDP can elicit  
11 transient bacterial adaptations, the mechanisms of which are different from those involved in the  
12 development of bacterial resistance to conventional antibiotics. For example, a study has shown that  
13 removal of CHDP from the culture medium resulted in the bacteria reverting to their original state: the  
14 adaptation to counter the effect of the CHDP was not maintained<sup>77</sup>. Therefore, microbial ‘resistance’,  
15 as described for conventional antibiotics, is unlikely to develop for CHDPs.

16 In addition to inducing gross damage of bacterial membranes, CHDP may also affect cell wall  
17 synthesis. For example, defensins such as HNP-1 and hBD3 exert their antibacterial activity by docking  
18 on lipid II, the intermediate in peptidoglycan synthesis<sup>78-80</sup>. In addition, CHDP such as the ribosome-  
19 binding proline-rich peptides can cross the bacterial membrane and kill bacteria from within, by  
20 binding to intracellular targets such as nucleic acids or nascent proteins and subsequently affect cell  
21 processes such as replication, transcription, translation, protein folding and cell wall synthesis<sup>28,76,81,82</sup>  
22 (Fig. 2). Simultaneous exposure of a pathogen to multiple different CHDP, potentially utilizing  
23 different modes of action, may be a critical mechanism by which these peptides are so effective *in vivo*.  
24 An exciting recent study in drosophila utilized CRISPR gene editing to delete all known immune-  
25 inducible CHDP and demonstrated both additive and synergistic antimicrobial functions of the  
26 peptides, as well as highly specific CHDP-pathogen interactions *in vivo*<sup>83</sup>.

27 It should be noted that the antibacterial effects of CHDP are measured under non-physiological  
28 conditions in most *in vitro* studies. This is a problem as the direct microbicidal activities of many  
29 CHDP are antagonized under physiological *in vivo* conditions, such as higher ionic strength, presence  
30 of divalent cations and host lipids and proteins. Thus, it may be argued that CHDP may not function *in*  
31 *vivo* as antimicrobials with direct microbicidal properties<sup>11,84</sup>. However, some CHDP (and derived  
32 synthetic peptides) with compromised direct antimicrobial activity *in vitro* exhibit the capacity to

actively control infections *in vivo*, which may be due to local concentrations of CHDP released by neutrophil degranulation at the site of infection being higher than *in vitro* MIC values. A more plausible explanation is that CHDP-mediated immunomodulatory functions and/or the concerted action of CHDP with other immunity-related factors are critical in the resolution of infections *in vivo* (discussed below). However, the relationship between antibacterial potency of CHDP *in vitro* and its immunomodulatory functions is not understood. There is no evidence of an inverse correlation between antibacterial potency and the ability of CHDP to induce an immune response<sup>84</sup>.

## **[H2] Antiviral Activity**

In addition to their antibacterial properties, early observations of the antiviral potential of CHDP<sup>85</sup> have been expanded to demonstrate activity against a range of viruses. The majority of these studies in recent years have been performed *in vitro* and have described a variety of mechanisms that underpin the antiviral effects, differing in the context of specific viruses (Table I).

A common mechanism of action against many enveloped viruses (such as Influenza, Respiratory Syncytial virus, Zika, Vaccinia virus, and Kaposi's sarcoma-associated herpesvirus) *in vitro* is the capacity of CHDP (including defensins and cathelicidins) to destabilise the viral envelope upon contact, damaging the virions and inhibiting infectivity<sup>86-94</sup>. This may happen upon contact in solution or upon viral exposure to plasma membrane-associated CHDP during cell entry<sup>88</sup>. However, CHDP have also been shown to have antiviral activity against non-enveloped viruses (such as Rhinovirus, Human Papillomavirus 16 and Adenovirus), by decreasing viral replication<sup>95,96</sup> and/or via binding viral capsid, thereby preventing uncoating and nuclear entry of the viral genome<sup>97-101</sup>. An additional mechanism of action against a number of viruses (such as Herpes Simplex Virus and HIV) relates to specific CHDP binding to cellular receptors involved in viral infection, that is dependent upon the lectin-like properties of some peptides<sup>102-110</sup>. Further antiviral effects may also result from CHDP-mediated aggregation of viral particles<sup>111</sup>, inhibition of PKC activity<sup>112,113</sup>, and immunomodulatory effects (discussed in detail below) of the peptides on host immune cells such as enhancing phagocyte function<sup>111,114</sup> or by modification of cytokine responses<sup>115,116</sup>. These antiviral mechanisms highlight the possibility that baseline expression of CHDP could create an “antiviral shield” at mucosal surfaces and prevent replication and spread of virus, if upregulated after initial infection. Therapies aimed at inducing host CHDP expression, or the application of synthetic CHDP-derived peptides with defined, selective properties, could therefore have both preventative and/or therapeutic potential.

## **[H2] Antifungal activity**

Fungal infections in humans are a growing problem world-wide. Many patients suffer from non-life-threatening fungal infections of the skin and oral cavity. However, invasive infections by fungal species such as *Aspergillus fumigatus*, *Cryptococcus neoformans*, *Candida albicans* and *Histoplasma capsulatum* are responsible for the death of 1.5 million people annually<sup>117</sup>. Only a limited set of antifungals is available for the treatment of life-threatening fungal infections and the development of antifungal drug resistance continues to increase<sup>118</sup>. This is particularly evident for azole-resistance because of the wide-spread use of azoles in agriculture<sup>119</sup>. In addition, the world-wide number of immunocompromised patients is increasing which leads to more fungal infections. Both trends are very disturbing and demand the development of new potent antifungals that do not easily elicit resistance. Furthermore, it is important that novel antifungal compounds be only used in humans to prevent rapid resistance development in the environment. Because of their diverse modes of action CHDP-derived molecules may be used as paradigms for the development of potent antifungals. Antifungal CHDP from plants and vertebrates have been described<sup>48</sup>. The activity of plant defensins has primarily been described against fungi<sup>120</sup>. Various killing mechanisms of CHDP on yeast and fungi have been reported, ranging from effects on mitochondrial functions of *Candida albicans* by histatin 5 to membrane effects on this yeast by cathelicidins and CHDP mimics<sup>121-123</sup>. It should be noted that fungal biofilms are highly resistant to antifungals<sup>124</sup>. It is thus important to screen for the anti-biofilm activity of newly developed CHDP-based antifungals<sup>124</sup>.

## **[H1] Immunomodulatory actions of CHDP**

The earliest studies into the non-microbicidal properties of CHDP were on their effects on immune cells, primarily related to the ability of these peptides to recruit leukocytes<sup>125-127</sup>. Following this, research on immunity-related functions of CHDP increased exponentially over the next two decades, defining a diverse range of functions. Immunity-related functions of CHDP seem to be dependent on the environmental stimuli, cell and tissue type, interaction with different cellular receptors, and the concentration of the peptides. Studies to date indicate that the molecular mechanism underpinning the ability of CHDP to selectively modulate immune responses is highly complex involving intracellular uptake of the peptides, which may or may not be mediated by membrane-associated GPCR, interaction with several intracellular interacting protein partners or receptors (e.g. GAPDH and p62), alteration of several signaling pathways (NFκB, p38 and JNK MAPK, MKP-1, and PI3K), engagement of different transcription factors, all of which seem to be dependent on factors such as the peptide concentration,

kinetic of response and the environmental stimuli (reviewed in<sup>12,15,128,129</sup>). The pleiotropic immunomodulatory functions of CHDP raise questions regarding the primary biological role of these peptides. Below, we summarize the activities of CHDP on modulation of immunity and inflammation (Fig. 3), primarily focusing on cathelicidins and defensins. Understanding mechanisms that underpin the ability of CHDP to modulate immunity to protect against infection, resolve inflammation and contribute to immune homeostasis, is critical to the development of novel therapeutic approaches based on CHDP-derived peptides.

## **[H2] Immune activation**

Protective activation of the innate immune system by CHDP has emerged as one of the key mechanisms underpinning the ability of these peptides to promote early clearance of infections. The actions of CHDP include recruiting leukocytes, modulating neutrophil responses and influencing antigen-specific adaptive immunity (discussed below). In addition, the interaction of CHDP and the microbiome in mucosal immunity is a rapidly emerging area of research. Although beyond the scope of this review, this exciting area of research suggests the potential for CHDP-mediated selectivity in control of the microbiome<sup>130-133</sup>, but also the potential for components of the microbiome to contribute to the regulation of CHDP expression. Further research on CHDP-microbiome interaction may enhance understanding of the consequences of microbial dysbiosis in disease states, aging and following treatment with broad-spectrum antibiotics.

## **[H3] Leukocyte recruitment**

Immune cells e.g. neutrophils, macrophages, mast cells and T-cells exhibit direct chemotactic activity towards CHDP and their IDR derivatives<sup>54,57,127,134-136</sup>. CHDP also indirectly promote recruitment of leukocytes by inducing the release of chemokines<sup>10,128,137-140</sup>. The ability of these peptides to induce chemokine release and enhance recruitment of leukocytes has been defined as a primary immunomodulatory mechanism related to their ability to protect against infections<sup>54,57,141</sup>. Underlying mechanisms involve several cellular receptors such as chemokine receptors (e.g. CCR6 and CCR2), G-protein coupled receptors (GPCR) including the formyl peptide receptors (reviewed in<sup>12,142</sup>), and Toll-like receptors (TLR)<sup>135,143,144</sup>, as well as interaction with intracellular proteins such as GAPDH and p62<sup>145,146</sup>. However, these functions appear to be dependent on the phase of infection and inflammation, as it has been shown that LL-37 can mediate the internalization of chemokine receptor CXCR2 on monocytes and neutrophils, consequently dampening chemotaxis<sup>147</sup>. Molecular processes

that control the dichotomy of CHDP to selectively induce chemokine secretion and enhance leukocyte recruitment, without altering the anti-inflammatory functions of the peptide (discussed in the next section) are not completely understood.

### *[H3] Neutrophil function*

Neutrophils are the major innate immune effector cells of the early phase response to infections, and are a primary source of both defensins and cathelicidins which are stored pre-formed in the secondary and primary granules respectively, and released upon neutrophil degranulation. CHDP can influence the function of neutrophils to modify infection outcomes. For example, these peptides can enhance influx of neutrophils both by direct chemotactic function<sup>127,136,144</sup> and indirectly by promoting the secretion of neutrophil chemokines e.g. IL-8 and Gro- $\alpha$  in a MAPK-dependent manner<sup>137,148</sup>, to promote control of infections. Similarly, a recent study has shown that CHDP hepcidin induced in the skin during bacterial infections can enhance chemokine production and promote neutrophil responses<sup>149</sup>. CHDP are also located on neutrophil NETs, potentially contributing to NET-mediated antibacterial effects<sup>150</sup>. LL-37 can facilitate NET formation and contribute to antiviral activity, as reported for influenza A virus<sup>86</sup>. It should be noted that recent studies have shown that post-translational modifications of CHDP, in particular citrullination, may alter the functions of NET associated CHDP such as cathelicidins<sup>151-153</sup>. LL-37 can also promote bacterial clearance *in vivo* by enhancing early neutrophil responses<sup>154</sup>. In addition, LL-37 can induce intracellular calcium mobilisation and the generation of reactive oxygen species (ROS)<sup>147</sup> as well as enhance ROS production mediated by other inflammatory stimuli<sup>155</sup>, indicating the capacity to prime and enhance neutrophil antimicrobial functions. The ability of CHDP to modulate the host cellular response to infections is not restricted to effects on neutrophils, these peptides can also modify other innate and adaptive cellular responses (reviewed in<sup>12,14,29,33</sup>).

### *[H3] Antigen presentation and adaptive immunity*

CHDP serve as a link between innate and adaptive immunity, as a consequence of their capacity to recruit antigen presenting cells (APCs) such as monocytes / macrophages and dendritic cells (DCs) to the site of infections (as discussed above). These peptides can also enhance phagocytosis in macrophages, facilitating clearance of bacteria by boosting immune activation<sup>156-159</sup>. CHDP can not only activate APCs, but also influence the generation and polarization of lymphocyte responses, thus shaping the adaptive immune response. For example, human defensins hBD2 and hBD3 can both

1 induce production of IFN- $\alpha$  from plasmacytoid DCs and consequently influence the initiation of T-cell  
2 responses<sup>160</sup>. The expression of co-stimulatory molecules CD80, CD86 and CD40 on myeloid cells are  
3 also upregulated by exposure to hBD3, facilitating the promotion of adaptive immune response via the  
4 engagement of TLRs<sup>161</sup>. LL-37 can modulate the adaptive immune response by multiple effector  
5 functions; LL-37 facilitates enhanced expression of co-stimulatory molecules and promotion of a  
6 modified adaptive response via modulation of dendritic cell differentiation and function *in vitro*<sup>159,162</sup>  
7 and *in vivo*<sup>163</sup>, promotes DC activation<sup>164,165</sup> and enhances the activation/ proliferation of B-cells via  
8 activating follicular DCs<sup>166</sup>. The influence of CHDP on adaptive immunity has led to the examination  
9 of application of these peptides, primarily cathelicidins, as adjuvants to enhance systemic and mucosal  
10 antigen-specific immune responses<sup>144,160,166</sup>.

11

## 12 **[H2] Regulation of inflammation**

13 As discussed above, CHDP exhibit multiple functions to activate the immune system which can be  
14 classified as pro-inflammatory responses that aid in the clearance of pathogens. In contrast, potent  
15 CHDP-mediated anti-inflammatory effects have been demonstrated in the presence of an inflammatory  
16 and/or pathogenic challenge, thus suggesting that CHDP are also regulatory molecules that limit  
17 enhanced inflammation. Therefore, rather than being described as either pro- or anti-inflammatory  
18 molecules, it may be appropriate to define CHDP as molecules that can balance inflammation to  
19 promote immune homeostasis. The anti-inflammatory function of CHDP is reinforced by several  
20 studies that have demonstrated that the deficiency of these peptides results in increased inflammatory  
21 responses; cathelicidin-deficient mice exhibit a more severe inflammatory phenotype compared to wild  
22 type<sup>155,167,168</sup>. Similarly, reduced expression of  $\beta$ -defensin in enterocytes of humans has been associated  
23 with Crohn's disease<sup>169</sup>. Indeed the critical role of defensins in maintaining the integrity of intestinal  
24 mucosa and immune homeostasis is well established (reviewed in<sup>34</sup>). Notably, exogenous application of  
25 CHDP e.g. LL-37, CATH-2 and BMAP-28, hBD2, and synthetic peptides e.g. IDR-1 and IDR-1002,  
26 has been shown to control inflammation in various animal models of infections and sepsis<sup>55,57,59,60,170-</sup>  
27 <sup>173</sup>. Similarly, an LL-37-derived peptide controlled the disease process in a murine model of  
28 inflammatory arthritis<sup>20</sup>, and IDR-1002 effectively alleviated airway inflammation *in vivo*<sup>21</sup>. In  
29 contrast, cathelicidin-KO mice demonstrate increased survival rate in a cecal ligation and puncture  
30 model of sepsis, despite increased expression of pro-inflammatory genes<sup>167</sup>. Therefore, the outcome of  
31 CHDP-mediated immunomodulatory responses seems to be context dependent and reliant on the

cellular environment. This has been demonstrated in studies examining the cross talk between TLR and CHDP, as discussed below.

### *[H3] Crosstalk of CHDP with TLRs*

Several studies have demonstrated that cathelicidins exhibit potent anti-endotoxin properties *in vitro* and *in vivo*<sup>10,138,170,174,175</sup>, both by binding bacterial lipopolysaccharide (LPS)<sup>174,175</sup> and by intervening in TLR signaling mechanisms<sup>138</sup>. The ability of cathelicidins such as LL-37 to modulate TLR-to-NFκB signaling is not restricted to its effect on TLR4, as LL-37 also dampens responses to TLR2/1 agonists and whole bacteria<sup>10,176</sup>. Downstream outcomes of CHDP-mediated modulation of the TLR-to-NFκB pathway in the presence of inflammatory stimuli results in the selective suppression of specific pro-inflammatory responses such as production of TNF and ROS, without the neutralization of innate immune functions such as chemokine production, and concurrent enhancement of anti-inflammatory mediators such as IL-10, IL-1RA, A20 and NFκBIA<sup>138,177-180</sup>. A caveat is that the effect of endogenous CHDP to limit inflammation may be impeded by peptide modifications that occur under inflammatory conditions. A recent study demonstrated that the enzyme peptidyl arginine deaminase released by inflammatory cells can mediate citrullination of LL-37, which compromises the ability of the peptide to intervene in TLR-signaling to dampen pro-inflammatory responses in macrophages, and to control sepsis *in vivo*<sup>151</sup>. Therefore, it is essential to consider the composition of the inflammatory milieu in studies that aim to define or optimize the anti-inflammatory functions of cationic peptides.

It should be noted that the modulation of TLR-mediated signalling by CHDP is not necessarily “anti-inflammatory”, as these peptides can also induce chemokines to attract leukocytes such as IL-8 and MCP-1 (pro-inflammatory response) at the same time as inhibiting LPS-induced TNF<sup>10,138,181</sup>. Similarly, cathelicidins can also exhibit both pro- or anti-inflammatory activities depending on the phase of the infection. For example, CATH-2 enhances the sensing of bacterial DNA<sup>182</sup> and is not anti-inflammatory if the bacterial infection has not been completely resolved<sup>182,183</sup>, while LL-37 promotes acute protective pro-inflammatory responses primarily in the context of an infection, including via activation of the inflammasome, but not in the absence of pathogens<sup>154,184</sup>. However, in instances where the bacteria are killed, both LL-37 and CATH-2 exhibit anti-inflammatory activity and can prevent activation via TLR2 and TLR4<sup>183</sup>. This is one of the mechanisms described for cathelicidin-mediated dampening of inflammatory responses induced by bacterial products, which may be especially important in pulmonary infections to protect the respiratory epithelium<sup>173</sup>.

Similarly, human defensins e.g. hBD3 exhibit anti-endotoxic properties *in vitro* and *in vivo*, via modulation of TLR signalling pathways<sup>185,186</sup>, but also induces the production of pro-inflammatory cytokines by monocytes<sup>161</sup>, increases TLR9-dependent responses to bacterial DNA<sup>187</sup>, and modifies MDA- and TLR-mediated responses to Poly I:C<sup>188</sup>. In addition, hBD3 can induce the maturation of dendritic cells (DC)<sup>161</sup>, while hBD2 and 3 can induce TLR9-dependent plasmacytoid DC IFN- $\alpha$  release, secondary to interaction between the peptides and otherwise non-inflammatory DNA to mediate *in vivo* adjuvant properties<sup>189</sup>. This is similar to the capacity to mediate TLR7-, TLR8- and TLR9-dependent immunogenic responses to self-DNA/RNA, which was first described for LL-37<sup>164,165,190</sup>. However, LL-37 enhances TLR3-mediated responses<sup>191,192</sup> and promotes release of IL-1 $\beta$ <sup>193</sup>. These observations demonstrate the potential of CHDP to modulate innate and downstream adaptive response via their impact upon pattern recognition signalling pathways.

### [H3] Modulation of cytokine-mediated responses

Limited studies have demonstrated the ability of CHDP, especially LL-37, to modulate cytokine-mediated responses in various cell types. LL-37 induces the expression of the IL-1 family of genes including Th17/Th1-related genes such as *IL-6* and *IL23A* in keratinocytes<sup>194</sup>. Consistent with this, another study demonstrated that LL-37 can synergize with cytokines IL-1 $\beta$  and GM-CSF to enhance chemokine production, but not with cytokines such as IL-4 or IL-12, in human PBMC<sup>195</sup>. In contrast, LL-37 suppresses cytokine IL-32-induced pro-inflammatory cytokines such as TNF, IL-6 and IL-1 $\beta$ , by activating the dual phosphatase MKP-1 which is a known negative regulator of inflammation, without altering its ability to induce chemokine production<sup>177</sup>. Mechanisms that control the ability of CHDP such as LL-37 to limit inflammation without altering chemokine production are not completely understood. To that end, a recent study has demonstrated that LL-37-induced chemokine release and subsequent leukocyte recruitment is selectively mediated by GPCR via the Cdc42 RhoGTPase pathway, without impacting LL-37-induced anti-inflammatory cytokine release<sup>137</sup>.

### [H2] Cell death

Although CHDP can rapidly permeabilize prokaryotic membranes, most natural peptides are relatively less toxic to eukaryotic cells. CHDP such as LL-37 and CATH-2 can enter eukaryotic cells<sup>182,196</sup> and facilitate the entry of nucleic acids and DNA dyes<sup>182,193,197</sup>, without inducing cell lysis, suggesting a temporary membrane disruption or pore opening. However, exposure to high concentrations of LL-37 induces eukaryotic cell apoptosis *in vitro* and *in vivo*<sup>198-201</sup>. This phenomenon is cell type-dependent



with the viability of primary human lymphocytes and monocytes relatively unaffected by exposure to high concentrations of the peptide<sup>159,202</sup>. The impact of CHDP-mediated induction of apoptosis on innate or adaptive responses *in vivo* remains unknown. However, high concentrations of LL-37 have been found to preferentially induce death in infected epithelial cells and represent a potential additional host defence mechanism<sup>184,199</sup>. Furthermore, different modes of cell death have important roles in maintaining immune homeostasis, and in amplifying or dampening, and later resolving, inflammatory responses. In this context, the control of neutrophil death and anti-inflammatory properties of apoptotic neutrophils is particularly important<sup>203</sup>. Interestingly, LL-37 can induce rapid secondary necrosis of apoptotic neutrophils<sup>178,204</sup>, with anti-inflammatory effects upon activated macrophages being associated with LL-37-mediated release of granule contents from the apoptotic cells, perhaps resulting from release of LL-37 and  $\alpha$ -defensins from these apoptotic neutrophils<sup>205</sup>. Thus, CHDP-mediated regulation of cell death has the potential to affect the magnitude and the resolution of inflammatory responses to infection.

#### **[H1] Development of CHDP-based therapies**

Based on the properties discussed above, many CHDP from either prokaryotes or eukaryotes and their derivatives are currently being investigated for a variety of indications including their use as antimicrobials and anti-inflammatories, as well as their application in wound healing,

#### **[H2] Antimicrobial therapies**

#### **[H3] Preclinical studies**

Studies in murine models have demonstrated the critical role of CHDP in the control of infections. For example, cathelicidin-deficient mice<sup>206</sup> exhibit less effective host defense against Streptococcal skin infections<sup>206</sup>, impaired clearance of *Pseudomonas aeruginosa* infections from the lung<sup>207</sup> and the cornea<sup>208</sup>, develop more severe pox skin lesions upon infection with Vaccinia virus<sup>92</sup>, develop more severe infection with RSV upon challenge<sup>90</sup>, and exhibit increased susceptibility to infection with *Citrobacter rodentium* in the intestinal tract<sup>209</sup>, *Escherichia coli* in the urinary tract<sup>210</sup> and *Klebsiella pneumoniae* in the lung<sup>211</sup>. The impaired antimicrobial activities observed in these cathelicidin-deficient mice may be explained by the absence of direct effects of the mouse cathelicidin, CRAMP, on these pathogens, although the observed effects may also be the result of altered immune modulation in the knock-out mice.

1 Exogenous administration of many CHDP has been found to be effective in various animal infection  
2 models for bacterial, viral, and fungal infections (reviewed in<sup>12,212</sup>). However, it remains to be  
3 determined if the observed efficacy is primarily due to the direct antimicrobial activities of the  
4 administered CHDP on the pathogen or due to the effect of the peptides on host immunity, or a  
5 combination of both. Therefore, the mode of action of the peptides needs to be carefully examined *in*  
6 *vivo*.

7 Several factors have limited the success of oral or systemic use of CHDP in preclinical studies,  
8 including (i) high local concentrations of CHDP, (ii) concerted actions of different CHDP and (iii)  
9 synergism with other molecules at the site of infection. However, many *in vivo* studies have shown  
10 efficacy of CHDP-derived peptides when these are administered topically. For example, a recent study  
11 showed that topical application of a peptide designed using LL-37 and Tachyplesin 1 as chemical  
12 benchmarks was protective in a MRSA murine model<sup>213</sup>. Similarly, localized intra-tracheal  
13 administration of a CHDP-derived peptide was shown to be better in lowering bacterial load in the  
14 lungs compared to rifampicin treatment in a murine model of tuberculosis<sup>214</sup>. Moreover, nanostructure-  
15 based technology has recently shown promise as an effective delivery system for slow release of  
16 peptides for infection control *in vivo*<sup>215</sup>.

17 A growing area of interest is the potential use of CHDP with indwelling medical devices, prosthetic  
18 joints and other implants for the prevention of nosocomial infections<sup>216,217</sup>. Bacterial biofilm formation  
19 on medical devices include pathogens recalcitrant to antibiotics which results in biomaterial-associated  
20 infections, a major problem in clinical practice. CHDP may be immobilized on the surfaces of  
21 biomaterials to prevent adhesion of bacteria. For example, synthetic peptides designed from LL-37 and  
22 a trombocidin-derived peptide was shown to be effective in inhibiting biofilm formation by a  
23 biomaterial-associated clinical isolate of *S. aureus*<sup>216</sup>. In addition, depending on the chemical tethering  
24 procedure, it is possible to retain antibacterial activity of CHDP after coating the surface. The  
25 disadvantage of this approach is that only bacteria in the immediate vicinity of the surface are killed.  
26 Application of CHDP-releasing biomaterials may be a better approach to prevent infections from  
27 implants. It has been reported that several hydrogels and also nanotubes and microporous calcium  
28 phosphate coatings inhibit bacterial growth *in vivo*<sup>217</sup>.

29 Another potential application of CHDP is their development as potential first line antiviral  
30 treatments for use during pandemics, where there is insufficient time to produce vaccines (such as new  
31 influenza (IAV) pandemics), or for viral infections for which vaccines are not available (such as  
32 Respiratory Syncytial Virus (RSV)), and more broadly for other viral pathogens. An early proof of

concept study in mice demonstrated that gel-based application of HD5 protected against HSV infection<sup>102</sup>. More recently, intravaginal instillation of a synthetic defensin, identified by HD5 mutant screening, showed prophylactic and/or therapeutic efficacy in a lethal HSV-2 infection model<sup>103</sup>. In another example, prophylactic application of RC-2 in a murine HSV-mediated ocular keratitis model modestly reduced viral titres and reduced disease<sup>105</sup>, but had no effect on disease pathology when applied post-infection. Intranasal human and murine cathelicidins both have shown *in vivo* antiviral activity equivalent to current first line neuraminidase inhibitors in a lethal murine IAV model, when applied concomitantly with virus and daily thereafter, dramatically improving survival despite modest effects on viral load<sup>116</sup>. Similarly, intranasal Urumin<sup>87</sup>, intranasal mBD2 (optimal when premixed with virus before infection<sup>218</sup>), intravenous delivery of recombinant mBD3<sup>219</sup> or intramuscular delivery of mBD1-mBD3 fusion genes<sup>220</sup> were all protective in murine lethal IAV infection models. These studies all demonstrate the therapeutic potential of CHDP as antivirals *in vivo*, although further investigation is needed. Many *in vivo* studies suggest that while early direct contact of virus and CHDP may be protective due to direct damage to the virions, later stage modulation of host immune / inflammatory responses by the peptides may also be critical in the antiviral activity of a CHDP<sup>115,221</sup>. Such observations highlight the importance of studies that address both the direct microbicidal activities and the immunomodulatory properties of these peptides.

### [H3] Clinical trials

Most CHDP in clinical trials so far have been formulated for topical applications or as inhalants for the treatment of infections (see <http://dramp.cpu-bioinform.org/browse/ClinicalTrialsData.php> and Table II<sup>222</sup>). One of the most advanced of these was Pexiganan, an analogue of the magainin peptide, which was tested as a topical cream for the treatment of infected diabetic foot ulcers in Phase III clinical trials. However, development was terminated because it did not perform better than current treatments. There are several trials in progress using Omiganan, a CHDP-derived antimicrobial compound (detailed in Table II). For example, a Phase III trial evaluating the long-term safety for topical application of Omiganan as a treatment for rosacea is ongoing. In addition, localized application of a human lactoferricin-derived peptide PXL01-containing hydrogel was shown to be safe, well tolerated and effective as anti-adhesion treatment postoperatively after tendon repair surgery, in an in-patient Phase II clinical trial<sup>223</sup>. Also, clinical trials are ongoing for topical application of LL-37 for treatment of venous leg ulcers (Table II)<sup>224</sup>.

A small number of clinical trials investigating the toxicity and efficacy of CHDP using oral and intravenous administration routes have also been conducted or are ongoing (Table II). For example, Isegaran, an analog of the peptide protegrin was used as an oral solution for oral mucositis in Phase III clinical trial, but did not show significant efficacy (Table II). Similarly, Surotomycin and Isegaran completed Phase III trials but were rejected for ongoing development either due to poor efficacy or efficacy that was not superior to current drugs (Table II). Phase III clinical studies using intravenous Brilacidin, a synthetic defensin mimetic), for skin infections are starting soon. Recently, Phase III trials of Murepavadin as an intravenous treatment for bacterial pneumonia were terminated due to increased serum creatinine levels in patients, indicative of acute kidney injury. This is reminiscent of nephrotoxicity issues with polymyxins, the cationic nonribosomal peptides which were used for treatment of Gram-negative bacterial infections and are currently used as the antibiotics of last resort.

Another strategy being evaluated and tested in clinical trials is to enhance the expression/production of endogenous CHDP for chronic inflammatory and infectious disease. A multicentre, double-blind, randomized clinical trial demonstrated that supplementation with vitamin D was beneficial to control exacerbations in COPD, especially for patients deficient in vitamin D<sup>225</sup>. Vitamin D<sub>3</sub> results in the induction of LL-37 in macrophages which has been also associated with intracellular killing of *Mycobacterium tuberculosis* in a human trial<sup>226</sup>. Similarly, oral phenylbutyrate, with or without vitamin D supplementation, leads to the induction of LL-37 in macrophages and lymphocytes, and has been evaluated in the treatment of adults with active pulmonary tuberculosis<sup>226</sup>. Therefore, strategies to enhance endogenous CHDP production may be valuable for antimicrobial therapies to counter challenges associated with peptide delivery, stability and bioavailability.

In summary, only a handful of peptide-derived treatments have made it to market, they include PAC-113, a histatin analog that is being sold in Taiwan as a topical treatment for oral candidiasis and Dalbavancin, a semisynthetic lipoglycopeptide that has been approved in the US for intravenous treatment of acute skin infections. Although the failure rate has so far been high, the number of CHDP in clinical trials has grown rapidly and is likely to lead to success in the future.

## **[H2] Immunomodulatory therapies**

### **[H3] Preclinical studies**

Early clinical trials using synthetic analogues of CHDP that had been designed to maximize microbicidal activity achieved only moderate efficacy<sup>227</sup>, perhaps due to failure to recognize the importance of immunity-related functions of these peptides. Despite issues of concentration at mucosal

surfaces and antagonizing factors at sites of inflammation, CHDP are clearly essential for the control of infections *in vivo*<sup>90,92,206-211</sup>. Application of LL-37 is protective against infection with *P. aeruginosa*, influenza and RSV *in vivo*<sup>89,116,228</sup>, with mode of action involving enhanced, protective early neutrophil responses, rather than by direct microbicidal activity against the pathogen<sup>154</sup>. Comparable results were also obtained after *in ovo* application of CATH-2 in chickens to induce a long-lasting protection against respiratory *E. coli* infections<sup>229</sup>. Similarly, CHDP-derived synthetic peptides such as IDR peptides were also shown to be protective in various infection models wherein IDR peptides protect the host from the pathogen by modulating the host immune response, and in parallel suppress the release of inflammatory cytokines such as TNF and IL-6, suppress reactive oxygen species (ROS), and dampen neutrophil degranulation<sup>54,57-59</sup>. These studies demonstrate that CHDP or related synthetic peptides can provide protection against infections by modulating host immune response rather than by directly targeting the pathogen.

Synthetic IDR peptides are primarily being developed as immunomodulatory therapies to control infections, with particular focus on antibiotic resistant infections. IDR-1 was the first such peptide shown to be protective against both Gram-positive and Gram-negative bacteria, including antibiotic-resistant infections such as methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus* (VRE) and *Salmonella* in murine preclinical models<sup>54</sup>. Subsequently, an analog of IDR-1, peptide SGX94, showed broad-spectrum activity against bacterial infections solely by modulating the host immune response to control infections in pre-clinical studies<sup>230</sup>. Other IDR peptides such as IDR-1002 and IDR-1018 have also been shown to be beneficial in preclinical models of a wide range of pathogens, including multidrug resistant bacteria, viruses, parasites, and antibiotic-recalcitrant bacterial biofilms<sup>56-59</sup>. In general, the mechanism of action of IDR peptides is primarily mediated by influencing host innate immune responses to enhance immune cell recruitment to the site of infection to promote bacterial clearance, rather than by directly targeting the pathogen. IDR peptides also influence neutrophil functions to augment neutrophil-mediated killing of bacteria and the release of natural CHDP from neutrophils<sup>231</sup>.

In addition to the potential application of CDHPs in infections, the immunomodulatory functions of these peptides are being examined for use in other indications. Cathelicidin-deficient mice develop more severe colitis in a non-infectious DSS-induced model with greater pro-inflammatory cytokine expression and cell death compared to wild type controls, and with impaired mucus production<sup>232</sup>. In contrast, therapeutic intrarectal administration of the murine cathelicidin mCRAMP (or mCRAMP-expressing *Lactococcus lactis*) is protective in the DSS-induced colitis murine model<sup>233,234</sup>. Similarly,

therapeutic administration of exogenous LL-37 or mCRAMP can modulate *C. difficile* colitis by inhibiting toxin-A-associated intestinal inflammation<sup>235</sup>. These studies suggest that CHDP and/or derived synthetic peptides have the potential to control chronic inflammatory diseases. Consistent with this, limited preclinical studies have demonstrated the beneficial effects of CHDP-derived synthetic peptides for chronic inflammatory diseases such as in inflammatory arthritis<sup>20</sup>, asthma<sup>21</sup>, and in colitis<sup>22</sup> models. For example, exogenous administration of a LL-37-derived synthetic peptide prevented the development of arthritis, suppressed autoantibodies and prevented cartilage degradation of the joints, in a collagen-induced arthritis murine model<sup>20</sup>. Elafin delivered by adenovirus ameliorated colitis, suppressed inflammatory cytokines and related NF- $\kappa$ B activation in non-infectious DSS-induced colitis murine model<sup>22</sup>. In addition, the peptide IDR-1002, a bactenecin derivative, improved allergen-induced airway hyperresponsiveness, controlled airway inflammation and suppressed inflammatory cytokine IL-33 production, in a murine model of asthma<sup>21</sup>. These preclinical studies suggest that CHDP and their derivative peptides can be developed for non-infectious inflammatory diseases.

Due to the ability of CHDP to regulate inflammation, many studies have also explored the effects of these peptides on cancers. However, the therapeutic use of CHDP-based peptides for cancers remains controversial, as the effect of these peptides on cancer pathology seems to be dependent on the specific type of cancer<sup>236</sup>. Discussing the nuances of pharmaceutical relevance of CHDP in cancer pathology is beyond the scope of this review<sup>237,238</sup>. Nevertheless, a gene therapy study using hDB2 showed that application of this peptide resulted in enhanced local anti-tumor effects in preclinical murine models using CT26, MethA and LL/2 tumor cells, the mechanism of which was associated with immunomodulatory functions of the peptide, namely its ability to activate endogenous dendritic cells<sup>23</sup>.

Another promising avenue for harnessing the immunomodulatory functions of IDR peptides is their application as adjuvants for new vaccine formulations, as IDR peptides enhance mucosal immunity and antigen-specific humoral responses<sup>61-64</sup>. Adjuvant formulations with IDR peptides have shown promise in various preclinical studies for vaccines against mycoplasma<sup>63</sup>, influenza H1N1 strain<sup>239</sup>, and pertussis<sup>240</sup>. In these studies, IDR peptides with modest antimicrobial activity used in different vaccine formulations showed profound immunomodulatory effects. For example, a bactenecin derived-IDR peptide used in a vaccine formulation with *M. bovis* proteins elicited a balanced humoral IgG1/IgG2 response for use in cattle<sup>63</sup>. Intranasal administration of a nanoparticle-based vaccine formulation containing an immunomodulatory IDR peptide as an adjuvant resulted in a strong cellular and humoral response against H1N1 influenza strain<sup>239</sup>. The potential use of IDR peptide as an adjuvant in vaccines is strongly supported by studies demonstrating that the presence of IDR peptide in these formulations is

effective in eliciting a balanced Th1/Th2 immune response along with facilitating the reduction of antigen dose required for immunity<sup>241</sup>.

The application of immunomodulatory CHDP in wound healing is also being explored, driven by studies showing that growth factors associated with stimulation of regeneration of tissues also induce the production of endogenous CHDP such as LL-37,  $\beta$ -defensin and lipocalin in keratinocytes<sup>242</sup>. Aligned with this, *in vitro* studies have demonstrated that CHDP such as defensins and LL-37 promote angiogenesis and wound healing<sup>243-246</sup>, prevent protease-mediated skin barrier damage<sup>247</sup> and promote re-epithelialisation of wounds<sup>248</sup>. These phenotypic changes are driven by peptide-mediated activation of signaling intermediates and transcription factors that activate EGFR through the induction of a G-protein-coupled receptor, in particular for the peptide LL-37<sup>244</sup>. Accordingly, application of LL-37 was shown to be effective in a murine model of excisional wound<sup>244</sup>.

### [H3] Clinical trials

Despite many preclinical studies describing the immunomodulatory therapeutic potential of CHDP-derived peptides, there have been very few successful clinical trials. The human cathelicidin LL-37 has been applied in a first-in-man randomized, placebo-controlled clinical trial, and shown to be both safe with no local or systemic adverse reactions and to enhance healing of hard-to-heal venous leg ulcers (Table II)<sup>224</sup>. A recent notable study in current clinical development is the use of silicone hydrogel contact lenses coated with a synthetic immunomodulatory peptide Mel4 to reduce contact lens-associated infections and inflammation<sup>249</sup>. Notably, the potential therapeutic and prophylactic success of immunomodulatory CHDP-derived peptides as alternatives to antibiotics in veterinary medicine may pave the way for clinical trials in humans (One Health approach)<sup>250</sup>.

### [H1] Considerations in CHDP-based drug development

The use of natural CHDP as effective therapeutics is not particularly viable as concentrations that exhibit direct antimicrobial effects are relatively high, and at that concentration range these peptides exhibit cytotoxic effects such as mast cell degranulation, complement activation and apoptosis of mammalian cells, and induce pro-inflammatory cytokine production<sup>11,12,15,251</sup>. Thus, synthetic peptides derived from natural CHDP, synthetic designed peptides and peptides found by semi-random high throughput screening are now emerging as putative lead compounds. Recent studies have also focused on non-peptide CHDP mimics such as peptoid analogs and developing compounds using CHDP on small abiotic scaffolds, for therapeutic applications (reviewed in<sup>24</sup>).

Major challenges facing peptide-based drug development include formulation and delivery, as well as high production cost. Biological factors that affect peptide stability and bioavailability must be taken into consideration, for example mucosal pH and the presence of host or microbial proteases that can degrade candidate peptides<sup>252-254</sup>, as well as several other factors which can impair peptide activity such as physiological salt concentration, mucus, DNA and microbial saccharides<sup>252,254</sup>. Several approaches are therefore being explored to enhance peptide stability, such as using D-amino acid peptides and modification of peptides by amidation or acetylation of the terminal regions or by targeted substitutions of tryptophan or histidine with a non-natural amino acid (reviewed in<sup>212</sup>).

The antimicrobial activity of CHDP is generally less than that of conventional antibiotics. Regulatory authorities require new antimicrobials to be non-inferior to existing antibiotics, even if the new compounds do not elicit antimicrobial resistance. CHDP may be unsuitable for standard *in vitro* antimicrobial susceptibility test methodologies to predict *in vivo* efficacy, even for topical application, and more physiologically relevant modified approaches may be vital<sup>255</sup>. Taking all this into consideration, the oral or systemic use of CHDP, with the aim to directly kill microorganisms, will likely be difficult to achieve. However, discoveries of CHDP demonstrating effects against antibiotic resistant pathogens do hold promise<sup>217</sup>, as does the new direction of applying immunomodulatory CHDP as an adjunct to antibiotics, due to the observed synergy of CHDP with conventional antibiotics<sup>256-259</sup>.

Additional approaches that are being considered to counter the challenges associated with CHDP-based therapy include the use of nutritional supplements such as vitamin D or phenylbutyrate or other products based on short chain fatty acids to enhance the levels of endogenous CHDP<sup>226</sup>. In addition, topical application of analogs of vitamin D was shown to enhance local expression of cathelicidin in psoriatic skin, which would circumvent any challenges that may be related to high systemic levels of vitamin D<sup>260,261</sup>. Also formulations using nanoparticles or liposomes for slow release and targeted delivery of CHDP<sup>262-264</sup>, as well as the use of shorter synthetic peptides such as IDR or cryptic peptides are being examined (discussed above). In preclinical studies<sup>56-59</sup>, shorter synthetic peptides such as IDR peptides have demonstrated negligible toxicity, no immunogenicity, and cost considerably less to produce than most CHDP, thus making these valuable candidates to investigate for clinical application. In addition, considerable less peptide is required if CHDP are used as immunomodulatory agents compared to their use as direct antimicrobial agents. Furthermore, new production methods such as the efficient usage of expression systems rather than chemical synthesis reduce the costs of production<sup>217,265,266</sup>.



Although CHDP offer a promising approach to treat infections, a key challenge for the use of CHDP or related synthetic analogues as antimicrobials is the development of pathogen-associated resistance mechanisms. Indeed resistance of pathogens to human CHDP has been demonstrated with multiple mechanisms adapted by bacteria to evade the direct antimicrobial effects of the peptides<sup>267-269</sup> (see Box. 1 and Fig. 4). However, bacteria have a limited number of ways to resist CHDP, and this resistance is often costly. An important recent study of *in vivo* survival and pathogenicity of a CHDP-resistant *S. aureus* (evolved *in vitro* in the presence of cationic peptides) showed that resistance to the peptides provided no survival advantage to the bacteria in an insect host environment that is dominated by antimicrobial peptides, bacterial clearance was at least as efficient as for the sensitive strains<sup>270</sup>. Nevertheless, harnessing the immunomodulatory actions of CHDP to selectively boost the host immune response rather than directly targeting the pathogen may be the path forward in the development of CHDP-based anti-infective therapies.

## [H1] Outlook

The repertoire of functions exhibited by CHDP ranges from direct antimicrobial activity to a wide range of effects on host defense mechanisms, highlighting the critical role of these molecules in infection and immunity. Although, research in this field was initially focused on the development of new ‘antibiotics’ based on cationic antimicrobial peptides, it is now well appreciated that CHDP have a critical role in immunity; from activation of innate immunity, enhancement of antigen presentation and phagocytosis, to influencing adaptive immunity and memory functions, along with potent anti-inflammatory functions. It is thus not surprising that research in this field has intensified in the context of drug development for a variety of clinical applications ranging from the control of antibiotic-resistant pathogens, alleviation of inflammation in chronic disease, their use as antibiotic adjuvants, to the targeting of specific cancers. However, there are many challenges associated with CHDP-based drug development, notably those associated with formulation and delivery, the potential for drug resistance, as well as the lack of solid pharmacokinetic data. Nevertheless, the wide range of CHDP functions defined to date provides a diverse range of natural molecules for the design and optimization of new drugs. Despite many associated challenges and the limited understanding of structure-function relationships, the potential of CHDP-based therapies remains a promising new clinical direction.

## Boxes

### ***Box 1: Development of antimicrobial resistance to CHDP:***

A major consideration for the potential application of CHDP as new generation antibiotics will have to include a thorough understanding of the frequency of resistance development of pathogens to peptide-based therapies. CHDP have promise over small molecule antibiotics because the surfaces with which they interact with targets in the pathogen are larger<sup>271</sup> and hence single amino acid substitutions are unlikely to lead to adaptations of bacteria to mitigate CHDP activity. Furthermore, CHDP have complicated mechanisms of action, often interacting with more than one target in microbes, such that multiple mutations within the pathogen are needed for ‘resistance’ to the peptides. Indeed, a recent study showed that bacterial adaptations to resist CHDP action do not develop easily<sup>272</sup>. The additional, indirect CHDP-mediated effects of enhancing host immune responses to control infections provide an important complementation to the direct microbicidal activities, providing a multi-faceted attack on pathogens during infection. Nevertheless, recent studies have revealed that bacterial and fungal pathogens are capable of developing mechanisms to resist the effects of CHDP. The mechanisms of adaptation of pathogens to human CHDP have been studied extensively and reviewed<sup>267-269</sup>. Common mechanisms in bacteria to counter the effects of CHDP are repulsion, sequestration, removal and degradation (Fig. 4). Additional mechanisms of bacterial adaptations are modification of the pentapeptide on Lipid II, a prominent CHDP target, and by altering the rigidity of the membrane by acylation of Lipid A. Mechanisms that fungi employ to enhance tolerance to CHDP have mainly focused on *Candida* species, which also employ repulsion, sequestration, removal by efflux pumps and proteolytic degradation against peptides such as LL-37, histatin 5, HNP-1, hBD-3 and lactoferrin<sup>273</sup>. Commensal bacteria of the host microbiome must be able to survive the CHDP presented by epithelial and mucosal surfaces. This may in part be due to the relatively low concentrations of CHDP at mucosal surfaces (except for specific niches such as the intestinal crypts) and somewhat inhibitory environments in which secreted CHDP are present normally in the absence of inflammation. In addition, proteases that cleave endogenous peptides to generate the active form of mature CHDP are also either absent or inactive in the absence of an inflammatory response. In contrast, CHDP in the phagolysosome of a neutrophil, for example, are at a high concentration in a controlled environment optimized for pathogen killing. Relative bacterial resistance to host CHDP is a prerequisite for effective commensal colonisation<sup>131</sup>, a property which may be most critical for stability through periods of inflammation where increased levels of CHDP may be capable of preferentially removing pathogens without totally decimating the healthy microbiome. Harnessing this sort of selectivity would have clear therapeutic advantages over broad spectrum antibiotics, and is indeed the focus of an exciting new approach to treating atopic dermatitis<sup>274</sup>, now being developed in human clinical trials.

## Figure legends

### Figure 1: Structures of CHDP

Part a illustrates examples of structures of CHDP from vertebrates. Two cathelicidins are depicted: human LL-37 and chicken CATH-2 (with proline-induced kink). Two human defensins are shown: the  $\alpha$ -defensin HD-5 and the  $\beta$ -defensin HBD-2. The pairing and positioning of the six conserved cysteine residues is as follows:  $\alpha$ -defensins: Cys1-Cys6, Cys2-Cys4, Cys3-Cys5;  $\beta$ -defensins: Cys1-Cys5, Cys2-Cys4, Cys3-Cys6. Magainin-2, a peptide from *Xenopus laevis*; its analogue, pexiganan, was developed as a topical agent. Amino acid side chains: red, hydrophobic; blue, basic; green, acidic. Part b provides examples of the diversity of CHDP structures. Tertiary structures of selected peptides from plants, fungi and invertebrates, arranged by secondary structure content. Beta strands shown in blue, alpha helices in red, and disulphide bonds in yellow (PDBs: 1MR4, 1NB1, 2RNG, 1BHP, 1GD3, 2L2R, 1JBL, 1HEV, 2MAL, 5E5Q, 5OQS).

**Figure 2: Models of antibacterial mechanisms of CHDP.** Direct antimicrobial mechanisms of CHDP can be mediated by membrane translocation of the peptides followed by binding to intracellular targets such as nucleic acids and/or proteins to kill bacteria. The mechanisms of translocation are not clear and may depend on the peptide and bacterial species. Proline-rich antimicrobial peptides use inner membrane transporters as Trojan horses to gain entry and subsequently bind within the ribosomal exit tunnel. Other CHDP may use transient pores for translocation. Interaction of CHDP with negatively charged bacterial membrane resulting in membrane perturbation has been defined as a primary mode of direct antimicrobial action. The models of membrane perturbation proposed are barrel-stave, carpet and toroidal pore models. The barrel-stave model was the first permeabilization mechanism proposed and considered to be the prototype in peptide-mediated transmembrane pore formation. In this model, peptides act as staves and vertically insert into the lipid bilayer forming barrels. Peptides which act according to the carpet model cover the negatively charged membrane based on electrostatic attraction. Above a certain peptide threshold concentration, the membrane ruptures in a detergent-like manner resulting in micelle formation of peptide with membrane lipids. The toroidal model is a variation of the aggregate model, where after parallel binding of the peptide to the membrane, the peptide distorts the alignment of the polar head groups of the lipids. This results in perturbation of the acyl chain interactions of the lipids, changes in membrane curvature and destabilization of membrane surface

integrity. At certain peptide to lipid ratios, the peptides orient perpendicularly to the membrane and induce the formation of transient toroidal channels.

**Figure 3: Summary of immunomodulatory mechanisms of CHDP.** Immunomodulatory functions exhibited by CHDPs include but are not limited to; recruitment of antigen presenting cells to site of infections either directly or indirectly by induction of chemokines to enhance antimicrobial effects, facilitating the activation of NETs, altering endotoxin-mediated signaling pathways, suppression of pro-inflammatory cytokines, enhancing phagocytosis and pro-inflammatory responses to nucleic acids, induction of anti-inflammatory cytokines, influencing differentiation of dendritic cell and polarization of T-cells. Adapted from van der Does A, Hiemstra P and Mookherjee N, *Adv Exp Med Biol* 2019<sup>15</sup>.

**Figure 4: Diagram of common resistance mechanisms to CHDP in bacterial and fungal pathogens.** (A) Gram positive bacteria (B) Gram negative bacteria (C) yeast/fungi. 1. Degradation by secreted proteases, outer membrane proteases or cytosolic proteases. 2. Sequestration by secreted proteins, anionic polysaccharides, mannosylphosphate side chains on glycoproteins (fungi) or O-antigen (Gram negative bacteria). 3. Electrostatic repulsion by alanylated lipoteichoic acid (LTA) or wall teichoic acid (WTA). 4. Electrostatic repulsion by aminoacylated phosphatidyl glycerol (PG). 5. Blocking CHDP binding by altering the pentapeptide on Lipid II. 6. Export of CHDP by efflux pumps. 7. Activation of signal transduction pathways that induce expression of genes that reinforce the wall or detoxify products of CHDP activity. 8. Lipid A modification by amine compounds. 9. Enhanced membrane rigidity by lipid A acylation. 10. Activation of MAPK signaling pathways in fungi for protection against oxidative, osmotic or cell wall stress. Adapted from Joo et al., 2016, and Swidergall and Ernst, 2014.

# 1 Tables

## 2 Table I: Antiviral activities of cationic host defence antimicrobial peptides

Virus	Peptide	Proposed mechanism of action <i>in vitro</i>	References
Influenza	HNP	<ul style="list-style-type: none"> <li>• Virus aggregation</li> <li>• Inhibition of PKC disrupts IAV endosomal trafficking</li> <li>• Enhanced neutrophil phagocytosis of IAV</li> </ul>	85,111,112
	Retrocyclins	<ul style="list-style-type: none"> <li>• Virus aggregation</li> <li>• Increased virus uptake by professional phagocytes</li> <li>• RC2: haemagglutinin-mediated fusion of viral and endosomal membranes blocked</li> </ul>	113,275,276
	$\beta$ -defensins	<ul style="list-style-type: none"> <li>• Inhibition of IAV infectivity at higher concentrations applied before viral entry</li> </ul>	114,115,219
	LL-37	<ul style="list-style-type: none"> <li>• Disruption of viral envelope</li> </ul>	116,218,277
	Urumin	<ul style="list-style-type: none"> <li>• Virion destruction, targeting H1 hemagglutinin</li> </ul>	87
RSV	hBD2	<ul style="list-style-type: none"> <li>• Viral envelope destabilisation in solution or upon exposure to plasma membrane-associated hBD2</li> </ul>	88,278
	LL-37	<ul style="list-style-type: none"> <li>• Virion binding and destruction</li> <li>• Prevention of infection and spread</li> <li>• Function retained by core 22-mer</li> </ul>	89,90
Rhinovirus	Cathelicidins	<ul style="list-style-type: none"> <li>• Decreased infectivity and replication</li> </ul>	91,95
Adenovirus	$\alpha$ -defensins	<ul style="list-style-type: none"> <li>• Peptide binding to adenoviral capsid prevents uncoating and nuclear entry of the viral genome</li> <li>• Dependent upon optimal peptide hydrophobicity and charge</li> </ul>	97-99,279
HPV 16	$\alpha$ -defensins	<ul style="list-style-type: none"> <li>• Uncoating and nuclear entry of the viral genome inhibited</li> </ul>	101
HSV	$\alpha$ -defensins, HBD3, retrocyclins	<ul style="list-style-type: none"> <li>• HSV binding to cellular receptors glycoprotein B and heparin sulphate inhibited</li> <li>• Dependent upon lectin-like properties,</li> </ul>	100,102,103,105

		rather than charge	
HIV	HNP	<ul style="list-style-type: none"> <li>• Disruption of cellular entry</li> <li>• Inhibition of PKC activity, interfering with HIV replication</li> </ul>	106,280-282
	Retrocyclins	<ul style="list-style-type: none"> <li>• Viral entry into cells blocked by peptide binding to gp120 and CD4</li> <li>• Dependent upon lectin-like properties</li> </ul>	107-110,283-285
	$\beta$ -defensins	<ul style="list-style-type: none"> <li>• Direct effects on virions</li> <li>• Intracellular, post-viral entry inhibitory functions</li> </ul>	286-288
	LL-37	<ul style="list-style-type: none"> <li>• Suppression of HIV reverse transcriptase activity</li> </ul>	289,290
Vaccinia virus	Cathelicidins	<ul style="list-style-type: none"> <li>• Integrity of the double layered viral envelope damaged</li> </ul>	92,291
Zika virus	Cathelicidins	<ul style="list-style-type: none"> <li>• Direct inactivation of Zika virus</li> <li>• Protective modulation of interferon signalling pathways</li> </ul>	93
Kaposi's sarcoma-associated herpesvirus		<ul style="list-style-type: none"> <li>• Disruption of viral envelope</li> </ul>	292

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12

1 **Table 2. Antimicrobial peptides under clinical trials.**

Peptide	Origin	Indication	Status	Company	Clinical trial identifier
<b>Topical</b>					
Pexiganan (Locilex®, MSI-78)	Analog of magainin, isolated from African clawed frog <i>Xenopus laevis</i>	Infected diabetic foot ulcers	Phase III complete. <i>Rejected, efficacy not superior</i>	PLx Pharma Inc. (formally Dipexium Pharmaceuticals Inc.)	<a href="#">NCT00563394</a> , <a href="#">NCT00563433</a> , <a href="#">NCT01590758</a> , <a href="#">NCT01594762</a>
D2A21, Demegel	Synthetic cecropin peptide	Burn wound infections	Phase III	Demegen	Not listed
CLS001 (Omiganan, MBI-226)	Omiganan pentahydrochloride. Synthetic cationic indolicidin derivative.	Local catheter site infections	Phase III complete (discontinued)	Mallinckrodt, BioWest Therapeutics Inc, Cadence Pharmaceuticals, Inc.	<a href="#">NCT00231153</a> , <a href="#">NCT00027248</a> , <a href="#">2005-003194-24</a>
		Topical skin antisepsis	Phase III complete	Mallinckrodt	<a href="#">NCT00608959</a>
		Papulopustular rosacea	Phase III current	Cutanea Life Sciences, Inc.	<a href="#">NCT02576860</a> , <a href="#">NCT02547441</a> , <a href="#">NCT02576847</a> , <a href="#">2015-002921-20</a> , <a href="#">2015-002919-15</a> , <a href="#">2015-002920-23</a>
		Acne vulgaris	Phase II complete	Cutanea Life Sciences, Inc., BioWest Therapeutics Inc	<a href="#">NCT02571998</a> , <a href="#">NCT02066545</a> , <a href="#">NCT00211497</a> , <a href="#">NCT00211523</a>
		Atopic dermatitis	Phase II complete	Cutanea Life Sciences, Inc.	<a href="#">NCT03091426</a> , <a href="#">NCT02456480</a> , <a href="#">2016-003849-28</a> , <a href="#">2014-003689-26</a>
		Vulvar intraepithelial neoplasia	Phase II complete	Cutanea Life Sciences, Inc.	<a href="#">NCT02596074</a> , <a href="#">2015-002724-16</a>
		Condylomata acuminata (external genital warts)	Phase II complete	Cutanea Life Sciences, Inc.	<a href="#">NCT02849262</a> , <a href="#">2015-005553-13</a>
		Facial seborrheic dermatitis	Phase II current	Cutanea Life Sciences, Inc./Maruho Co., Ltd	<a href="#">NCT03688971</a> , <a href="#">2017-003106-41</a>
Iseganan (IB-367)	Analog of protegrin-1	Ventilator-associated pneumonia	Phase II/III. <i>Rejected, no efficacy</i>	IntraBiotics Pharmaceuticals	<a href="#">NCT00118781</a>
PXL01	Synthetic macrocyclic 25 amino acid peptide derived from human lactoferricin	Prevention of post-surgical adhesions and scar prevention	Phase IIb complete.  Phase III trials planned	Promore Pharma (formally Pergamum AB)	<a href="#">NCT01022242</a> , <a href="#">2009-012703-25</a>

NVXT (Novexatin® NP213)	Cyclic arginine- based heptamer	Fungal nail infection (onychomycosis)	Phase IIb complete	NovaBiotics	<a href="#">NCT02933879</a> , <a href="#">NCT02343627</a>
PAC-113, P- 113	Histatin 5 derivative (12 amino acids)	Oral candidiasis	Phase IIb complete	Demegen/Pacgen Biopharmaceutical s Co. <b>Sold over the counter in Taiwan by General Biologicals Corporation</b>	<a href="#">NCT00659971</a>
LL-37	Human cathelicidin subunit	Venous leg ulcers	Phase IIb current	Promore Pharma (formally Pergamum AB)	<a href="#">2018-000536-10</a>
HXP124	Plant defensin	Fungal nail infection (onychomycosis)	Phase IIa complete	Hexima	<a href="#">ACTRN126180001 31257</a>
Brilacidin (PMX-30063)	Synthetic defensin mimetic	Ulcerative proctitis / ulcerative proctosigmoiditis	Phase II complete. Phase III planned	Alfasigma S.p.A	Not listed
		Oral mucositis in patients with head and neck cancer	Phase II complete  FDA Fast track designation	Innovation Pharmaceuticals (formally Cellceutix)	<a href="#">NCT02324335</a> , <a href="#">NCT01211470</a>
LTX-109 (Lytixar™)	Synthetic cationic tripeptide	Atopic dermatitis, skin infection	Phase II complete	Lytix Biopharma	<a href="#">NCT01223222</a> , <a href="#">2010-021438-68</a>
		Impetigo	Phase II complete	Lytix Biopharma	<a href="#">NCT01803035</a>
		Nasal infections by methicillin- resistant/-sensitive <i>Staphylococcus aureus</i> (MRSA/MSSA)	Phase I/II complete	Lytix Biopharma	<a href="#">NCT01158235</a> , <a href="#">2010-019254-40</a>
(CKPV)2, CZEN-002	Derivative of $\alpha$ - melanocyte stimulating hormone	Vulvovaginal candidiasis	Phase II complete	Zengen/Abiogen Pharma	<a href="#">2005-001360-31</a>
OP-145 (AMP60.4Ac)	Cathelicidin family (LL-37 derivative)	Chronic suppurative otitis media (middle ear infections)	Phase II complete	OctoPlus BV/Dr Reddy's Research and Development BV	<a href="#">ISRCTN1214972 0</a>
C16G2	Synthetic peptide	Prevention of dental caries due to <i>Streptococcus mutans</i>	Phase II complete	Armata Pharmaceuticals	<a href="#">NCT03052842</a> , <a href="#">NCT03004365</a> , <a href="#">NCT02594254</a> , <a href="#">NCT02509845</a> , <a href="#">NCT02254993</a> , <a href="#">NCT02044081</a> , <a href="#">NCT03196219</a>
DPK 060	Derived from kininogen, cationic random coil peptide	Acute external otitis	Phase II complete	DermaGen AB and Promore Pharma (formally Pergamum AB)	<a href="#">NCT01447017</a> , <a href="#">2011-004356-20</a>



		Atopic dermatitis	Phase I/II complete	DermaGen AB and Promore Pharma (formally Pergamum AB)	<a href="#">NCT01522391</a>
<a href="#">Lotilibcin</a> (WAP-8294A <sub>2</sub> )	Lipodepsipeptide	Methicillin-resistant <i>S. aureus</i> (MRSA)	Phase I complete	aRigen Pharmaceuticals/ Green Cross Corporation	Not listed
PL-5	Alpha helical peptide	Bacterial skin infections	Approval by State Food and Drug Administration of China (SFDA) for clinical trial	Changchun ProteLight Pharmaceutical & Biotechnology Co	n/a
<b>Oral</b>					
Surotomicin (CB-183, 315)	Cyclic lipopeptide, analog of daptamycin	Diarrhea caused by <i>Clostridium difficile</i>	Phase III complete. <b><i>Rejected, efficacy not superior</i></b>	Cubist Pharmaceuticals/Merk & Co., Inc.	<a href="#">NCT01597505</a> , <a href="#">NCT01598311</a> , <a href="#">2012-000252-3</a>
Iseganan (IB-367)	Analog of protegrin-1	Oral mucositis in patients with head and neck cancer	Phase III complete. <b><i>No efficacy</i></b>	National Cancer Institute (NCI)/ IntraBiotics Pharmaceuticals	<a href="#">NCT00022373</a>
RDP58, Delmitide acetate, allotrap 1258	d-amino acid decapeptide	Ulcerative colitis	Phase II complete	Genzyme/Procter & Gamble	<a href="#">2004-004077-29</a>
NVB-302	Synthetic type B lantibiotic	<i>Clostridium difficile</i> infection	Phase I complete	Novacta	<a href="#">ISRCTN40071144</a>
<b>Intravenous</b>					
Dalbavancin (BI397, Dalvance®, Xydalba™)	Semisynthetic lipoglycopeptide	Acute bacterial skin infections	Approved	Allergan (formally Actavis and Durata Therapeutics)	n/a
		Osteomyelitis and septic arthritis	Phase IV current		<a href="#">NCT03426761</a>
AB103 (p2TA)	Synthetic anionic CD28 dimer mimetic peptide	Necrotizing soft tissue infections	Phase III current	Atox Bio Ltd	<a href="#">NCT02469857</a> , <a href="#">2018-001125-15</a>
Dusquetide (SGX942)	Synthetic 5 amino acid peptide derived from indolizidine, immunomodulator	Oral mucositis in patients with head and neck cancer	Phase III current. FDA Fast track designation	Soligenix	<a href="#">NCT03237325</a> , <a href="#">2017-003702-41</a>
Murepavadin (POL7080)	Synthetic cyclic beta hairpin peptidomimetic based on the cationic antimicrobial peptide protegrin I	Ventilator-associated bacterial pneumonia by <i>Pseudomonas aeruginosa</i>	Phase III <b><i>suspended, adverse events</i></b>	Polyphor Ltd	<a href="#">NCT03409679</a> , <a href="#">NCT03582007</a>
Neuprex®, opebacan, bactericidal/permeability-	BPI-derived peptide	Burns	Phase II complete	University of Texas Southwestern Medical Center	<a href="#">NCT00462904</a>

increasing protein rBPI <sub>21</sub>		Myeloablative allogeneic hematopoietic stem cell transplantation	Phase I/II. <b>Terminated, lack of enrollment</b>	Xoma LLC	<a href="#">NCT00454155</a>
Brilacidin (PMX-30063)	Synthetic defensin mimetic	Acute Bacterial Skin and Skin Structure Infections (ABSSSI)	Phase II complete. Phase III planned FDA Fast track designation	Innovation Pharmaceuticals (formally Cellceutix)	<a href="#">NCT02052388</a>
EA-360	Linear tetrapeptide, derived from human chorionic gonadotropin hormone (hCG)	Systemic inflammatory response and renal function	Phase IIa/b current	Exponential Biotherapies	<a href="#">NCT03145220</a> , <a href="#">2014-002481-78</a>
hLF1-11 (human lactoferrin 1-11)	First cationic domain of human lactoferrin hLF (11 residues)	Infections during haematopoietic stem cell transplantations	Phase I/II complete. <b>withdrawn</b>	AM-Pharma	<a href="#">NCT00509938</a> , <a href="#">NCT00430469</a>
		Candidaemia	Phase I/II <b>withdrawn</b>	AM-Pharma	<a href="#">NCT00509834</a>
		Bacteremia due to Staphylococcus epidermidis	Phase I/II <b>withdrawn</b>	AM-Pharma	<a href="#">NCT00509847</a>
Friulimicin B	Cyclic lipopeptide	Pneumonia, Staphylococcal skin infections	Phase I. <b>Rejected, unfavourable pharmacokinetics</b>	MerLion Pharmaceuticals	<a href="#">NCT00492271</a>

## 1 Reference cited

- 2 1 Iannella, H., Luna, C. & Waterer, G. Inhaled corticosteroids and the increased risk of pneumonia: what's  
3 new? A 2015 updated review. *Ther Adv Respir Dis* **10**, 235-255, doi:10.1177/1753465816630208  
4 (2016).
- 5 2 Widdifield, J. *et al.* Serious infections in a population-based cohort of 86,039 seniors with rheumatoid  
6 arthritis. *Arthritis Care Res (Hoboken)* **65**, 353-361, doi:10.1002/acr.21812 (2013).
- 7 3 Clancy, C. J. *et al.* Emerging and resistant infections. *Ann Am Thorac Soc* **11 Suppl 4**, S193-200,  
8 doi:10.1513/AnnalsATS.201402-069PL (2014).
- 9 4 Simmaco, M., Kreil, G. & Barra, D. Bombinins, antimicrobial peptides from *Bombina* species. *Biochim*  
10 *Biophys Acta* **1788**, 1551-1555, doi:10.1016/j.bbamem.2009.01.004 (2009).
- 11 5 Steiner, H., Hultmark, D., Engstrom, A., Bennich, H. & Boman, H. G. Sequence and specificity of two  
12 antibacterial proteins involved in insect immunity. *Nature* **292**, 246-248 (1981).
- 13 6 Ganz, T. *et al.* Defensins. Natural peptide antibiotics of human neutrophils. *J Clin Invest* **76**, 1427-1435,  
14 doi:10.1172/JCI112120 (1985).
- 15 7 Zasloff, M. Magainins, a class of antimicrobial peptides from *Xenopus* skin: isolation, characterization of  
16 two active forms, and partial cDNA sequence of a precursor. *Proc Natl Acad Sci U S A* **84**, 5449-5453  
17 (1987).
- 18 8 Bucki, R., Byfield, F. J. & Janmey, P. A. Release of the antimicrobial peptide LL-37 from DNA/F-actin  
19 bundles in cystic fibrosis sputum. *Eur Respir J* **29**, 624-632, doi:10.1183/09031936.00080806 (2007).
- 20 9 Bergsson, G. *et al.* LL-37 complexation with glycosaminoglycans in cystic fibrosis lungs inhibits  
21 antimicrobial activity, which can be restored by hypertonic saline. *J Immunol* **183**, 543-551,  
22 doi:10.4049/jimmunol.0803959 (2009).
- 23 **10 Scott, M. G., Davidson, D. J., Gold, M. R., Bowdish, D. & Hancock, R. E. The human antimicrobial peptide**  
24 **LL-37 is a multifunctional modulator of innate immune responses. *J Immunol* **169**, 3883-3891 (2002).**  
25 **One of the first study to demonstrate the immunomodulatory properties of CHDP LL-37 and discuss**  
26 **immunomodulation as being critical to the antimicrobial functions of the peptide.**
- 27 11 Bowdish, D. M., Davidson, D. J., Scott, M. G. & Hancock, R. E. Immunomodulatory activities of small  
28 host defense peptides. *Antimicrob Agents Chemother* **49**, 1727-1732, doi:10.1128/AAC.49.5.1727-  
29 1732.2005 (2005).
- 30 12 Hancock, R. E., Haney, E. F. & Gill, E. E. The immunology of host defence peptides: beyond antimicrobial  
31 activity. *Nat Rev Immunol* **16**, 321-334, doi:10.1038/nri.2016.29 (2016).
- 32 13 Pantic, J. M. *et al.* The Potential of Frog Skin-Derived Peptides for Development into Therapeutically-  
33 Valuable Immunomodulatory Agents. *Molecules* **22**, doi:10.3390/molecules22122071 (2017).
- 34 14 Hemshekhar, M., Anaparti, V. & Mookherjee, N. Functions of Cationic Host Defense Peptides in  
35 Immunity. *Pharmaceuticals (Basel)* **9**, doi:10.3390/ph9030040 (2016).
- 36 15 van der Does, A. M., Hiemstra, P. S. & Mookherjee, N. Antimicrobial Host Defence Peptides:  
37 Immunomodulatory Functions and Translational Prospects. *Adv Exp Med Biol* **1117**, 149-171,  
38 doi:10.1007/978-981-13-3588-4\_10 (2019).
- 39 16 Mant, C. T. *et al.* De Novo Designed Amphipathic alpha-Helical Antimicrobial Peptides Incorporating  
40 Dab and Dap Residues on the Polar Face To Treat the Gram-Negative Pathogen, *Acinetobacter*  
41 *baumannii*. *J Med Chem* **62**, 3354-3366, doi:10.1021/acs.jmedchem.8b01785 (2019).
- 42 17 Jiang, S., Deslouches, B., Chen, C., Di, M. E. & Di, Y. P. Antibacterial Properties and Efficacy of a Novel  
43 SPLUNC1-Derived Antimicrobial Peptide, alpha4-Short, in a Murine Model of Respiratory Infection.  
44 *MBio* **10**, doi:10.1128/mBio.00226-19 (2019).
- 45 18 Mishra, B., Reiling, S., Zarena, D. & Wang, G. Host defense antimicrobial peptides as antibiotics: design  
46 and application strategies. *Curr Opin Chem Biol* **38**, 87-96, doi:10.1016/j.cbpa.2017.03.014 (2017).
- 47 19 Mai, S. *et al.* Potential applications of antimicrobial peptides and their mimics in combating caries and  
48 pulpal infections. *Acta Biomater* **49**, 16-35, doi:10.1016/j.actbio.2016.11.026 (2017).

1 20 Chow, L. N. *et al.* Human cathelicidin LL-37-derived peptide IG-19 confers protection in a murine model  
2 of collagen-induced arthritis. *Mol Immunol* **57**, 86-92, doi:S0161-5890(13)00493-8 [pii]  
3 10.1016/j.molimm.2013.08.011 (2013).

4 **21** Piyadasa, H. *et al.* Immunomodulatory innate defence regulator (IDR) peptide alleviates airway  
5 inflammation and hyper-responsiveness. *Thorax*, doi:10.1136/thoraxjnl-2017-210739 (2018). **The first**  
6 **study to demonstrate the therapeutic potential of an IDR peptide in allergen-challenged airway**  
7 **inflammation in the context of asthma.**

8 22 Ho, S., Pothoulakis, C. & Koon, H. W. Antimicrobial peptides and colitis. *Curr Pharm Des* **19**, 40-47  
9 (2013).

10 23 Li, D. *et al.* Gene therapy with beta-defensin 2 induces antitumor immunity and enhances local  
11 antitumor effects. *Hum Gene Ther* **25**, 63-72, doi:10.1089/hum.2013.161 (2014).

12 24 Scott, R. W. & Tew, G. N. Mimics of Host Defense Proteins; Strategies for Translation to Therapeutic  
13 Applications. *Curr Top Med Chem* **17**, 576-589, doi:10.2174/1568026616666160713130452 (2017).

14 25 Wang, G., Li, X. & Wang, Z. APD3: the antimicrobial peptide database as a tool for research and  
15 education. *Nucleic Acids Res* **44**, D1087-1093, doi:10.1093/nar/gkv1278 (2016).

16 26 Murakami, M., Lopez-Garcia, B., Braff, M., Dorschner, R. A. & Gallo, R. L. Postsecretory processing  
17 generates multiple cathelicidins for enhanced topical antimicrobial defense. *J Immunol* **172**, 3070-3077  
18 (2004).

19 **27** Sochacki, K. A., Barns, K. J., Bucki, R. & Weisshaar, J. C. Real-time attack on single *Escherichia coli* cells  
20 by the human antimicrobial peptide LL-37. *Proc Natl Acad Sci U S A* **108**, E77-  
21 81, doi:10.1073/pnas.1101130108 (2011). **The real-time attack on *E.coli* by the cathelicidin LL-37 is**  
22 **visualized for the first time.**

23 28 Schneider, V. A. *et al.* Imaging the antimicrobial mechanism(s) of cathelicidin-2. *Sci Rep* **6**, 32948,  
24 doi:10.1038/srep32948 (2016).

25 29 van Harten, R. M., van Woudenberg, E., van Dijk, A. & Haagsman, H. P. Cathelicidins:  
26 Immunomodulatory Antimicrobials. *Vaccines (Basel)* **6**, doi:10.3390/vaccines6030063 (2018).

27 30 Mookherjee, N., Rehaume, L. M. & Hancock, R. E. Cathelicidins and functional analogues as antiseptics  
28 molecules. *Expert Opin Ther Targets* **11**, 993-1004, doi:10.1517/14728222.11.8.993 (2007).

29 31 Zanetti, M. Cathelicidins, multifunctional peptides of the innate immunity. *J Leukoc Biol* **75**, 39-48,  
30 doi:10.1189/jlb.0403147  
31 jlb.0403147 [pii] (2004).

32 32 Patil, A., Hughes, A. L. & Zhang, G. Rapid evolution and diversification of mammalian alpha-defensins as  
33 revealed by comparative analysis of rodent and primate genes. *Physiol Genomics* **20**, 1-11,  
34 doi:10.1152/physiolgenomics.00150.2004 (2004).

35 33 Ganz, T. Defensins: antimicrobial peptides of innate immunity. *Nat Rev Immunol* **3**, 710-720,  
36 doi:10.1038/nri1180 (2003).

37 34 Bevins, C. L. & Salzman, N. H. Paneth cells, antimicrobial peptides and maintenance of intestinal  
38 homeostasis. *Nat Rev Microbiol* **9**, 356-368, doi:nrmicro2546 [pii]  
39 10.1038/nrmicro2546 (2011).

40 35 Semple, F. & Dorin, J. R. beta-Defensins: multifunctional modulators of infection, inflammation and  
41 more? *J Innate Immun* **4**, 337-348, doi:000336619 [pii]  
42 10.1159/000336619 (2012).

43 36 Bevins, C. L., Jones, D. E., Dutra, A., Schaffzin, J. & Muenke, M. Human enteric defensin genes:  
44 chromosomal map position and a model for possible evolutionary relationships. *Genomics* **31**, 95-106,  
45 doi:10.1006/geno.1996.0014 (1996).

46 37 Semple, C. A., Rolfe, M. & Dorin, J. R. Duplication and selection in the evolution of primate beta-  
47 defensin genes. *Genome Biol* **4**, R31, doi:10.1186/gb-2003-4-5-r31 (2003).

1 38 Morrison, G. M., Semple, C. A., Kilanowski, F. M., Hill, R. E. & Dorin, J. R. Signal sequence conservation  
2 and mature peptide divergence within subgroups of the murine beta-defensin gene family. *Mol Biol*  
3 *Evol* **20**, 460-470, doi:10.1093/molbev/msg060 (2003).

4 39 Gudmundsson, G. H. *et al.* The human gene FALL39 and processing of the cathelin precursor to the  
5 antibacterial peptide LL-37 in granulocytes. *Eur J Biochem* **238**, 325-332 (1996). **A seminal study which**  
6 **defines post-translational processing of cathelin precursor pro-peptide to the mature CHDP LL-37.**

7 40 van Dijk, A. *et al.* Chicken heterophils are recruited to the site of Salmonella infection and release  
8 antibacterial mature Cathelicidin-2 upon stimulation with LPS. *Mol Immunol* **46**, 1517-1526,  
9 doi:10.1016/j.molimm.2008.12.015 (2009).

10 41 Gallo, R. L. *et al.* Identification of CRAMP, a cathelin-related antimicrobial peptide expressed in the  
11 embryonic and adult mouse. *J Biol Chem* **272**, 13088-13093 (1997).

12 42 Melino, S., Santone, C., Di Nardo, P. & Sarkar, B. Histatins: salivary peptides with copper(II)- and zinc(II)-  
13 binding motifs: perspectives for biomedical applications. *FEBS J* **281**, 657-672, doi:10.1111/febs.12612  
14 (2014).

15 43 Goldstein, E. J. C., Citron, D. M., Tyrrell, K. L. & Leoncio, E. S. In Vitro Activity of Pexiganan and 10  
16 Comparator Antimicrobials against 234 Isolates, Including 93 Pasteurella Species and 50 Anaerobic  
17 Bacterial Isolates Recovered from Animal Bite Wounds. *Antimicrob Agents Chemother* **61**,  
18 doi:10.1128/AAC.00246-17 (2017).

19 44 Mylonakis, E., Podsiadlowski, L., Muhammed, M. & Vilcinskas, A. Diversity, evolution and medical  
20 applications of insect antimicrobial peptides. *Philos Trans R Soc Lond B Biol Sci* **371**,  
21 doi:10.1098/rstb.2015.0290 (2016).

22 45 Destoumieux-Garzon, D. *et al.* Antimicrobial peptides in marine invertebrate health and disease. *Philos*  
23 *Trans R Soc Lond B Biol Sci* **371**, doi:10.1098/rstb.2015.0300 (2016).

24 46 Shafee, T. M., Lay, F. T., Hulett, M. D. & Anderson, M. A. The Defensins Consist of Two Independent,  
25 Convergent Protein Superfamilies. *Mol Biol Evol* **33**, 2345-2356, doi:10.1093/molbev/msw106 (2016).

26 47 Shafee, T. M., Lay, F. T., Phan, T. K., Anderson, M. A. & Hulett, M. D. Convergent evolution of defensin  
27 sequence, structure and function. *Cell Mol Life Sci* **74**, 663-682, doi:10.1007/s00018-016-2344-5 (2017).

28 48 Vriens, K., Cammue, B. P. & Thevissen, K. Antifungal plant defensins: mechanisms of action and  
29 production. *Molecules* **19**, 12280-12303, doi:10.3390/molecules190812280 (2014).

30 49 Haney, E. F., Mansour, S. C., Hilchie, A. L., de la Fuente-Nunez, C. & Hancock, R. E. High throughput  
31 screening methods for assessing antibiofilm and immunomodulatory activities of synthetic peptides.  
32 *Peptides*, doi:S0196-9781(15)00073-X [pii]  
33 10.1016/j.peptides.2015.03.015 (2015).

34 50 de la Fuente-Nunez, C., Cardoso, M. H., de Souza Candido, E., Franco, O. L. & Hancock, R. E. Synthetic  
35 antibiofilm peptides. *Biochim Biophys Acta* **1858**, 1061-1069, doi:10.1016/j.bbamem.2015.12.015  
36 (2016).

37 51 Hilpert, K., Winkler, D. F. & Hancock, R. E. Peptide arrays on cellulose support: SPOT synthesis, a time  
38 and cost efficient method for synthesis of large numbers of peptides in a parallel and addressable  
39 fashion. *Nat Protoc* **2**, 1333-1349, doi:10.1038/nprot.2007.160 (2007).

40 52 Niyonsaba, F., Someya, A., Hirata, M., Ogawa, H. & Nagaoka, I. Evaluation of the effects of peptide  
41 antibiotics human beta-defensins-1/-2 and LL-37 on histamine release and prostaglandin D(2)  
42 production from mast cells. *Eur J Immunol* **31**, 1066-1075 (2001).

43 53 Chen, X. *et al.* Antimicrobial peptides human beta-defensin (hBD)-3 and hBD-4 activate mast cells and  
44 increase skin vascular permeability. *Eur J Immunol* **37**, 434-444, doi:10.1002/eji.200636379 (2007).

45 54 Scott, M. G. *et al.* An anti-infective peptide that selectively modulates the innate immune response.  
46 *Nat Biotechnol* **25**, 465-472, doi:nbt1288 [pii]  
47 10.1038/nbt1288 (2007). **The first study to demonstrate the ability of a synthetic IDR peptide in the control**  
48 **of bacterial infections including antibiotic-resistant MRSA and VRE in preclinical models.**

1 55 Rivas-Santiago, B. *et al.* Ability of Innate Defence Regulator Peptides IDR-1002, IDR-HH2 and IDR-1018  
2 to Protect against Mycobacterium tuberculosis Infections in Animal Models. *PLoS One* **8**, e59119,  
3 doi:10.1371/journal.pone.0059119

4 PONE-D-12-34871 [pii] (2013).

5 56 Achtman, A. H. *et al.* Effective adjunctive therapy by an innate defense regulatory Peptide in a  
6 preclinical model of severe malaria. *Sci Transl Med* **4**, 135ra164, doi:4/135/135ra64 [pii]  
7 10.1126/scitranslmed.3003515 (2012). **The first study to demonstrate effective adjuvant therapeutic  
8 potential of a IDR peptide with existing drugs in a parasitic infection model.**

9 57 Nijnik, A. *et al.* Synthetic cationic peptide IDR-1002 provides protection against bacterial infections  
10 through chemokine induction and enhanced leukocyte recruitment. *J Immunol* **184**, 2539-2550,  
11 doi:jimmunol.0901813 [pii]  
12 10.4049/jimmunol.0901813 (2010).

13 58 Mansour, S. C., de la Fuente-Nunez, C. & Hancock, R. E. Peptide IDR-1018: modulating the immune  
14 system and targeting bacterial biofilms to treat antibiotic-resistant bacterial infections. *J Pept Sci* **21**,  
15 323-329, doi:10.1002/psc.2708 (2015).

16 59 Wuerth, K. C., Falsafi, R. & Hancock, R. E. W. Synthetic host defense peptide IDR-1002 reduces  
17 inflammation in Pseudomonas aeruginosa lung infection. *PLoS One* **12**, e0187565,  
18 doi:10.1371/journal.pone.0187565 (2017).

19 60 Hou, M. *et al.* Antimicrobial peptide LL-37 and IDR-1 ameliorate MRSA pneumonia in vivo. *Cell Physiol  
20 Biochem* **32**, 614-623, doi:000354465 [pii]  
21 10.1159/000354465 (2013).

22 61 Cao, D. *et al.* CpG oligodeoxynucleotide synergizes innate defense regulator peptide for enhancing the  
23 systemic and mucosal immune responses to pseudorabies attenuated virus vaccine in piglets in vivo.  
24 *Int Immunopharmacol* **11**, 748-754, doi:S1567-5769(11)00063-4 [pii]  
25 10.1016/j.intimp.2011.01.028 (2011).

26 62 Kindrachuk, J. *et al.* A novel vaccine adjuvant comprised of a synthetic innate defence regulator peptide  
27 and CpG oligonucleotide links innate and adaptive immunity. *Vaccine* **27**, 4662-4671, doi:S0264-  
28 410X(09)00782-8 [pii]  
29 10.1016/j.vaccine.2009.05.094 (2009).

30 63 Prysliak, T. *et al.* Induction of a balanced IgG1/IgG2 immune response to an experimental challenge  
31 with Mycoplasma bovis antigens following a vaccine composed of Emulsigen, IDR peptide1002, and  
32 poly I:C. *Vaccine* **35**, 6604-6610, doi:10.1016/j.vaccine.2017.10.037 (2017).

33 64 Prysliak, T. & Perez-Casal, J. Immune responses to Mycoplasma bovis proteins formulated with  
34 different adjuvants. *Can J Microbiol* **62**, 492-504, doi:10.1139/cjm-2015-0762 (2016).

35 65 Wu, B. C., Lee, A. H. & Hancock, R. E. W. Mechanisms of the Innate Defense Regulator Peptide-1002  
36 Anti-Inflammatory Activity in a Sterile Inflammation Mouse Model. *J Immunol* **199**, 3592-3603,  
37 doi:10.4049/jimmunol.1700985 (2017).

38 66 Hoeksema, M., van Eijk, M., Haagsman, H. P. & Hartshorn, K. L. Histones as mediators of host defense,  
39 inflammation and thrombosis. *Future Microbiol* **11**, 441-453, doi:10.2217/fmb.15.151 (2016).

40 67 Urban, C. F. *et al.* Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex  
41 involved in host defense against Candida albicans. *PLoS Pathog* **5**, e1000639,  
42 doi:10.1371/journal.ppat.1000639 (2009).

43 68 Donovan, S. M. The Role of Lactoferrin in Gastrointestinal and Immune Development and Function: A  
44 Preclinical Perspective. *J Pediatr* **173 Suppl**, S16-28, doi:10.1016/j.jpeds.2016.02.072 (2016).



- 69 Bruni, N. *et al.* Antimicrobial Activity of Lactoferrin-Related Peptides and Applications in Human and Veterinary Medicine. *Molecules* **21**, doi:10.3390/molecules21060752 (2016).
- 70 Lehrer, R. I., Cole, A. M. & Selsted, M. E. theta-Defensins: cyclic peptides with endless potential. *J Biol Chem* **287**, 27014-27019, doi:R112.346098 [pii]
- 10.1074/jbc.R112.346098 (2012).
- 71 Forde, E. B. *et al.* Using Disease-Associated Enzymes to Activate Antimicrobial Peptide Prodrugs. *Methods Mol Biol* **1548**, 359-368, doi:10.1007/978-1-4939-6737-7\_26 (2017).
- 72 Pane, K. *et al.* Antimicrobial potency of cationic antimicrobial peptides can be predicted from their amino acid composition: Application to the detection of "cryptic" antimicrobial peptides. *J Theor Biol* **419**, 254-265, doi:10.1016/j.jtbi.2017.02.012 (2017).
- 73 Gaglione, R. *et al.* Novel human bioactive peptides identified in Apolipoprotein B: Evaluation of their therapeutic potential. *Biochem Pharmacol* **130**, 34-50, doi:10.1016/j.bcp.2017.01.009 (2017).
- 74 Tucker, A. T. *et al.* Discovery of Next-Generation Antimicrobials through Bacterial Self-Screening of Surface-Displayed Peptide Libraries. *Cell* **172**, 618-628 e613, doi:10.1016/j.cell.2017.12.009 (2018).
- 75 Torres, M. D. T., Sothiselvam, S., Lu, T. K. & de la Fuente-Nunez, C. Peptide Design Principles for Antimicrobial Applications. *J Mol Biol*, doi:10.1016/j.jmb.2018.12.015 (2019).
- 76 Jenssen, H., Hamill, P. & Hancock, R. E. Peptide antimicrobial agents. *Clin Microbiol Rev* **19**, 491-511 (2006).
- 77 Veldhuizen, E. J., Brouwer, E. C., Schneider, V. A. & Fluit, A. C. Chicken cathelicidins display antimicrobial activity against multiresistant bacteria without inducing strong resistance. *PLoS One* **8**, e61964, doi:10.1371/journal.pone.0061964 (2013).
- 78 Grein, F., Schneider, T. & Sahl, H. G. Docking on Lipid II-A Widespread Mechanism for Potent Bactericidal Activities of Antibiotic Peptides. *J Mol Biol* **431**, 3520-3530, doi:10.1016/j.jmb.2019.05.014 (2019).
- 79 de Leeuw, E. *et al.* Functional interaction of human neutrophil peptide-1 with the cell wall precursor lipid II. *FEBS Lett* **584**, 1543-1548, doi:10.1016/j.febslet.2010.03.004 (2010).
- 80 Sass, V. *et al.* Human beta-defensin 3 inhibits cell wall biosynthesis in Staphylococci. *Infect Immun* **78**, 2793-2800, doi:10.1128/IAI.00688-09 (2010).
- 81 Schneider, V. A. F. *et al.* Imaging the Antistaphylococcal Activity of CATH-2: Mechanism of Attack and Regulation of Inflammatory Response. *mSphere* **2**, doi:10.1128/mSphere.00370-17 (2017).
- 82 Graf, M. *et al.* Proline-rich antimicrobial peptides targeting protein synthesis. *Nat Prod Rep* **34**, 702-711, doi:10.1039/c7np00020k (2017).
- 83** Hanson, M. A. *et al.* Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *Elife* **8**, doi:10.7554/eLife.44341 (2019). **Using a systematic knockout approach, this study demonstrates that CHDP have synergistic, additive and highly specific microbicidal effects in vivo.**
- 84 Coorens, M., Scheenstra, M. R., Veldhuizen, E. J. & Haagsman, H. P. Interspecies cathelicidin comparison reveals divergence in antimicrobial activity, TLR modulation, chemokine induction and regulation of phagocytosis. *Sci Rep* **7**, 40874, doi:10.1038/srep40874 (2017).
- 85 Daher, K. A., Selsted, M. E. & Lehrer, R. I. Direct inactivation of viruses by human granulocyte defensins. *J Virol* **60**, 1068-1074 (1986).
- 86 Tripathi, S., Verma, A., Kim, E. J., White, M. R. & Hartshorn, K. L. LL-37 modulates human neutrophil responses to influenza A virus. *J Leukoc Biol* **96**, 931-938, doi:10.1189/jlb.4A1113-604RR (2014).
- 87 Holthausen, D. J. *et al.* An Amphibian Host Defense Peptide Is Virucidal for Human H1 Hemagglutinin-Bearing Influenza Viruses. *Immunity* **46**, 587-595, doi:10.1016/j.immuni.2017.03.018 (2017).
- 88 Kota, S. *et al.* Role of human beta-defensin-2 during tumor necrosis factor-alpha/NF-kappaB-mediated innate antiviral response against human respiratory syncytial virus. *J Biol Chem* **283**, 22417-22429, doi:10.1074/jbc.M710415200 (2008).

- Currie, S. M. *et al.* The human cathelicidin LL-37 has antiviral activity against respiratory syncytial virus. *PLoS One* **8**, e73659, doi:10.1371/journal.pone.0073659 (2013).
- Currie, S. M. *et al.* Cathelicidins Have Direct Antiviral Activity against Respiratory Syncytial Virus In Vitro and Protective Function In Vivo in Mice and Humans. *J Immunol* **196**, 2699-2710, doi:10.4049/jimmunol.1502478 (2016).
- Harcourt, J. L. *et al.* Human cathelicidin, LL-37, inhibits respiratory syncytial virus infection in polarized airway epithelial cells. *BMC Res Notes* **9**, 11, doi:10.1186/s13104-015-1836-y (2016).
- Howell, M. D. *et al.* Selective killing of vaccinia virus by LL-37: implications for eczema vaccinatum. *J Immunol* **172**, 1763-1767 (2004).
- Dean, R. E. *et al.* A carpet-based mechanism for direct antimicrobial peptide activity against vaccinia virus membranes. *Peptides* **31**, 1966-1972, doi:10.1016/j.peptides.2010.07.028 (2010).
- Brice, D. C., Toth, Z. & Diamond, G. LL-37 disrupts the Kaposi's sarcoma-associated herpesvirus envelope and inhibits infection in oral epithelial cells. *Antiviral Res* **158**, 25-33, doi:10.1016/j.antiviral.2018.07.025 (2018).
- Schogler, A. *et al.* Vitamin D represses rhinovirus replication in cystic fibrosis cells by inducing LL-37. *Eur Respir J* **47**, 520-530, doi:10.1183/13993003.00665-2015 (2016).
- Sousa, F. H. *et al.* Cathelicidins display conserved direct antiviral activity towards rhinovirus. *Peptides* **95**, 76-83, doi:10.1016/j.peptides.2017.07.013 (2017).
- Bastian, A. & Schafer, H. Human alpha-defensin 1 (HNP-1) inhibits adenoviral infection in vitro. *Regul Pept* **101**, 157-161 (2001).
- Smith, J. G. & Nemerow, G. R. Mechanism of adenovirus neutralization by Human alpha-defensins. *Cell Host Microbe* **3**, 11-19, doi:10.1016/j.chom.2007.12.001 (2008).
- Smith, J. G. *et al.* Insight into the mechanisms of adenovirus capsid disassembly from studies of defensin neutralization. *PLoS Pathog* **6**, e1000959, doi:10.1371/journal.ppat.1000959 (2010).
- Tenge, V. R., Gounder, A. P., Wiens, M. E., Lu, W. & Smith, J. G. Delineation of interfaces on human alpha-defensins critical for human adenovirus and human papillomavirus inhibition. *PLoS Pathog* **10**, e1004360, doi:10.1371/journal.ppat.1004360 (2014).
- Nguyen, E. K., Nemerow, G. R. & Smith, J. G. Direct evidence from single-cell analysis that human {alpha}-defensins block adenovirus uncoating to neutralize infection. *J Virol* **84**, 4041-4049, doi:10.1128/JVI.02471-09 (2010).
- Hazrati, E. *et al.* Human alpha- and beta-defensins block multiple steps in herpes simplex virus infection. *J Immunol* **177**, 8658-8666 (2006).
- Wang, A. *et al.* Enhancement of antiviral activity of human alpha-defensin 5 against herpes simplex virus 2 by arginine mutagenesis at adaptive evolution sites. *J Virol* **87**, 2835-2845, doi:10.1128/JVI.02209-12 (2013).
- Yasin, B. *et al.* Theta defensins protect cells from infection by herpes simplex virus by inhibiting viral adhesion and entry. *J Virol* **78**, 5147-5156 (2004).
- Brandt, C. R. *et al.* Evaluation of a theta-defensin in a Murine model of herpes simplex virus type 1 keratitis. *Invest Ophthalmol Vis Sci* **48**, 5118-5124, doi:10.1167/iovs.07-0302 (2007).
- Lehrer, R. I. *et al.* Multivalent binding of carbohydrates by the human alpha-defensin, HD5. *J Immunol* **183**, 480-490, doi:10.4049/jimmunol.0900244 (2009).
- Cole, A. M. *et al.* Retrocyclin: a primate peptide that protects cells from infection by T- and M-tropic strains of HIV-1. *Proc Natl Acad Sci U S A* **99**, 1813-1818, doi:10.1073/pnas.052706399 (2002).
- Gallo, S. A. *et al.* Theta-defensins prevent HIV-1 Env-mediated fusion by binding gp41 and blocking 6-helix bundle formation. *J Biol Chem* **281**, 18787-18792, doi:10.1074/jbc.M602422200 (2006).
- Wang, W. *et al.* Activity of alpha- and theta-defensins against primary isolates of HIV-1. *J Immunol* **173**, 515-520 (2004).
- Furci, L., Sironi, F., Tolazzi, M., Vassena, L. & Lusso, P. Alpha-defensins block the early steps of HIV-1 infection: interference with the binding of gp120 to CD4. *Blood* **109**, 2928-2935, doi:10.1182/blood-2006-05-024489 (2007).



1 111 Tecle, T., White, M. R., Gantz, D., Crouch, E. C. & Hartshorn, K. L. Human neutrophil defensins increase  
2 neutrophil uptake of influenza A virus and bacteria and modify virus-induced respiratory burst  
3 responses. *J Immunol* **178**, 8046-8052 (2007).

4 112 Salvatore, M. *et al.* alpha-Defensin inhibits influenza virus replication by cell-mediated mechanism(s). *J*  
5 *Infect Dis* **196**, 835-843, doi:10.1086/521027 (2007).

6 113 Sieczkarski, S. B., Brown, H. A. & Whittaker, G. R. Role of protein kinase C beta11 in influenza virus entry  
7 via late endosomes. *J Virol* **77**, 460-469 (2003).

8 114 Doss, M. *et al.* Hapivirins and diprovirins: novel theta-defensin analogs with potent activity against  
9 influenza A virus. *J Immunol* **188**, 2759-2768, doi:10.4049/jimmunol.1101335 (2012).

10 115 Ryan, L. K. *et al.* Modulation of human beta-defensin-1 (hBD-1) in plasmacytoid dendritic cells (PDC),  
11 monocytes, and epithelial cells by influenza virus, Herpes simplex virus, and Sendai virus and its  
12 possible role in innate immunity. *J Leukoc Biol* **90**, 343-356, doi:10.1189/jlb.0209079 (2011).

13 116 Barlow, P. G. *et al.* Antiviral activity and increased host defense against influenza infection elicited by  
14 the human cathelicidin LL-37. *PLoS One* **6**, e25333, doi:10.1371/journal.pone.0025333 (2011). **This**  
15 **study demonstrates the antiviral activity of CHDP against influenza virus *in vitro* and *in vivo*.**

16 117 Kim, J. Y. Human fungal pathogens: Why should we learn? *J Microbiol* **54**, 145-148,  
17 doi:10.1007/s12275-016-0647-8 (2016).

18 118 Revie, N. M., Iyer, K. R., Robbins, N. & Cowen, L. E. Antifungal drug resistance: evolution, mechanisms  
19 and impact. *Curr Opin Microbiol* **45**, 70-76, doi:10.1016/j.mib.2018.02.005 (2018).

20 119 Verweij, P. E., Chowdhary, A., Melchers, W. J. & Meis, J. F. Azole Resistance in *Aspergillus fumigatus*:  
21 Can We Retain the Clinical Use of Mold-Active Antifungal Azoles? *Clin Infect Dis* **62**, 362-368,  
22 doi:10.1093/cid/civ885 (2016).

23 120 Parisi, K. *et al.* The evolution, function and mechanisms of action for plant defensins. *Semin Cell Dev*  
24 *Biol* **88**, 107-118, doi:10.1016/j.semcdb.2018.02.004 (2019).

25 121 Menzel, L. P. *et al.* Potent *in vitro* and *in vivo* antifungal activity of a small molecule host defense  
26 peptide mimic through a membrane-active mechanism. *Sci Rep* **7**, 4353, doi:10.1038/s41598-017-  
27 04462-6 (2017).

28 122 Ordonez, S. R., Amarullah, I. H., Wubbolts, R. W., Veldhuizen, E. J. & Haagsman, H. P. Fungicidal  
29 mechanisms of cathelicidins LL-37 and CATH-2 revealed by live-cell imaging. *Antimicrob Agents*  
30 *Chemother* **58**, 2240-2248, doi:10.1128/AAC.01670-13 (2014).

31 123 Puri, S. & Edgerton, M. How does it kill?: understanding the candidacidal mechanism of salivary histatin  
32 5. *Eukaryot Cell* **13**, 958-964, doi:10.1128/EC.00095-14 (2014).

33 124 Delattin, N., Brucker, K., Cremer, K., Cammue, B. P. & Thevissen, K. Antimicrobial Peptides as a Strategy  
34 to Combat Fungal Biofilms. *Curr Top Med Chem* **17**, 604-612 (2017). **This study discusses recent**  
35 **advances using novel CHDP-based peptides to combat fungal biofilms.**

36 125 Chertov, O. *et al.* Identification of defensin-1, defensin-2, and CAP37/azurocidin as T-cell  
37 chemoattractant proteins released from interleukin-8-stimulated neutrophils. *J Biol Chem* **271**, 2935-  
38 2940 (1996).

39 126 Van Wetering, S. *et al.* Effect of defensins on interleukin-8 synthesis in airway epithelial cells. *Am J*  
40 *Physiol* **272**, L888-896, doi:10.1152/ajplung.1997.272.5.L888 (1997).

41 127 Yang, D., Chen, Q., Chertov, O. & Oppenheim, J. J. Human neutrophil defensins selectively  
42 chemoattract naive T and immature dendritic cells. *J Leukoc Biol* **68**, 9-14 (2000).

43 128 Choi, K. Y. & Mookherjee, N. Multiple immune-modulatory functions of cathelicidin host defense  
44 peptides. *Front Immunol* **3**, 149, doi:10.3389/fimmu.2012.00149 (2012).

45 129 Steinstraesser, L., Kraneburg, U., Jacobsen, F. & Al-Benna, S. Host defense peptides and their  
46 antimicrobial-immunomodulatory duality. *Immunobiology* **216**, 322-333, doi:S0171-2985(10)00136-1  
47 [pii]  
48 10.1016/j.imbio.2010.07.003 (2011).

1 130 Salzman, N. H. *et al.* Enteric defensins are essential regulators of intestinal microbial ecology. *Nat*  
2 *Immunol* **11**, 76-83, doi:10.1038/ni.1825 (2010).

3 131 Cullen, T. W. *et al.* Gut microbiota. Antimicrobial peptide resistance mediates resilience of prominent  
4 gut commensals during inflammation. *Science* **347**, 170-175, doi:10.1126/science.1260580 (2015).

5 132 Yoshimura, T. *et al.* The Antimicrobial Peptide CRAMP Is Essential for Colon Homeostasis by  
6 Maintaining Microbiota Balance. *J Immunol* **200**, 2174-2185, doi:10.4049/jimmunol.1602073 (2018).

7 133 Lee, R. J. *et al.* Bitter and sweet taste receptors regulate human upper respiratory innate immunity. *J*  
8 *Clin Invest* **124**, 1393-1405, doi:10.1172/JCI72094 (2014).

9 134 Madera, L. & Hancock, R. E. Synthetic immunomodulatory peptide IDR-1002 enhances monocyte  
10 migration and adhesion on fibronectin. *J Innate Immun* **4**, 553-568, doi:10.1159/000338648 (2012).

11 135 De, Y. *et al.* LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl  
12 peptide receptor-like 1 (FPRL1) as a receptor to chemoattract human peripheral blood neutrophils,  
13 monocytes, and T cells. *J Exp Med* **192**, 1069-1074 (2000).

14 136 Tjabringa, G. S., Ninaber, D. K., Drijfhout, J. W., Rabe, K. F. & Hiemstra, P. S. Human cathelicidin LL-37 is  
15 a chemoattractant for eosinophils and neutrophils that acts via formyl-peptide receptors. *Int Arch*  
16 *Allergy Immunol* **140**, 103-112 (2006).

17 137 Hemshekhar, M., Choi, K. G. & Mookherjee, N. Host Defense Peptide LL-37-Mediated Chemoattractant  
18 Properties, but Not Anti-Inflammatory Cytokine IL-1RA Production, Is Selectively Controlled by Cdc42  
19 Rho GTPase via G Protein-Coupled Receptors and JNK Mitogen-Activated Protein Kinase. *Front*  
20 *Immunol* **9**, 1871, doi:10.3389/fimmu.2018.01871 (2018).

21 138 Mookherjee, N. *et al.* Modulation of the TLR-mediated inflammatory response by the endogenous  
22 human host defense peptide LL-37. *J Immunol* **176**, 2455-2464, doi:10.1176/4/2455 [pii] (2006).

23 139 Yang, D. *et al.* LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl  
24 peptide receptor-like 1 (FPRL1) as a receptor to chemoattract human peripheral blood neutrophils,  
25 monocytes, and T cells. *J. Exp. Med* **192**, 1069-1074 (2000).

26 140 Tjabringa, G. S., Ninaber, D. K., Drijfhout, J. W., Rabe, K. F. & Hiemstra, P. S. Human cathelicidin LL-37 is  
27 a chemoattractant for eosinophils and neutrophils that acts via formyl-peptide receptors. *Int. Arch.*  
28 *Allergy Immunol* **140**, 103-112 (2006).

29 141 Holly, M. K., Diaz, K. & Smith, J. G. Defensins in Viral Infection and Pathogenesis. *Annu Rev Virol* **4**, 369-  
30 391, doi:10.1146/annurev-virology-101416-041734 (2017).

31 142 Agier, J., Efenberger, M. & Brzezinska-Blaszczyk, E. Cathelicidin impact on inflammatory cells. *Cent Eur J*  
32 *Immunol* **40**, 225-235, doi:10.5114/ceji.2015.51359 (2015).

33 143 Suarez-Carmona, M., Hubert, P., Delvenne, P. & Herfs, M. Defensins: "Simple" antimicrobial peptides or  
34 broad-spectrum molecules? *Cytokine Growth Factor Rev* **26**, 361-370, doi:S1359-6101(14)00169-5 [pii]  
35 10.1016/j.cytogfr.2014.12.005 (2015).

36 144 Kurosaka, K., Chen, Q., Yarovsky, F., Oppenheim, J. J. & Yang, D. Mouse cathelin-related antimicrobial  
37 peptide chemoattracts leukocytes using formyl peptide receptor-like 1/mouse formyl peptide receptor-  
38 like 2 as the receptor and acts as an immune adjuvant. *J Immunol* **174**, 6257-6265, doi:10.1176/4/6257  
39 [pii] (2005).

40 145 Mookherjee, N. *et al.* Intracellular receptor for human host defense peptide LL-37 in monocytes. *J*  
41 *Immunol* **183**, 2688-2696, doi:jimmunol.0802586 [pii]  
42 10.4049/jimmunol.0802586 (2009).

43 146 Yu, H. B. *et al.* Sequestosome-1/p62 is the key intracellular target of innate defense regulator peptide. *J*  
44 *Biol Chem* **284**, 36007-36011, doi:C109.073627 [pii]  
45 10.1074/jbc.C109.073627 (2009).

46 147 Zhang, Z. *et al.* Evidence that cathelicidin peptide LL-37 may act as a functional ligand for CXCR2 on  
47 human neutrophils. *Eur J Immunol* **39**, 3181-3194, doi:10.1002/eji.200939496 (2009).

1 148 Zheng, Y. *et al.* Cathelicidin LL-37 induces the generation of reactive oxygen species and release of  
2 human alpha-defensins from neutrophils. *Br J Dermatol* **157**, 1124-1131, doi:10.1111/j.1365-  
3 2133.2007.08196.x (2007).

4 149 Malerba, M. *et al.* Epidermal hepcidin is required for neutrophil response to bacterial infection. *J Clin*  
5 *Invest*, doi:10.1172/JCI126645 (2019).

6 150 Stephan, A. *et al.* LL37:DNA complexes provide antimicrobial activity against intracellular bacteria in  
7 human macrophages. *Immunology* **148**, 420-432, doi:10.1111/imm.12620 (2016).

8 151 Koziel, J. *et al.* Citrullination alters immunomodulatory function of LL-37 essential for prevention of  
9 endotoxin-induced sepsis. *J Immunol* **192**, 5363-5372, doi:10.4049/jimmunol.1303062 (2014).

10 152 Wong, A. *et al.* A Novel Biological Role for Peptidyl-Arginine Deiminases: Citrullination of Cathelicidin  
11 LL-37 Controls the Immunostimulatory Potential of Cell-Free DNA. *J Immunol* **200**, 2327-2340,  
12 doi:10.4049/jimmunol.1701391 (2018).

13 153 Kilsgard, O. *et al.* Peptidylarginine deiminases present in the airways during tobacco smoking and  
14 inflammation can citrullinate the host defense peptide LL-37, resulting in altered activities. *Am J Respir*  
15 *Cell Mol Biol* **46**, 240-248, doi:10.1165/rcmb.2010-0500OC (2012).

16 **154** Beaumont, P. E. *et al.* Cathelicidin host defence peptide augments clearance of pulmonary  
17 *Pseudomonas aeruginosa* infection by its influence on neutrophil function in vivo. *PLoS One* **9**, e99029,  
18 doi:10.1371/journal.pone.0099029 (2014). **This study demonstrates that cathelicidins can promote**  
19 **bacterial clearance in vivo, in the absence of direct microbicidal effects, by modulating host cellular**  
20 **inflammatory responses.**

21 155 Alalwani, S. M. *et al.* The antimicrobial peptide LL-37 modulates the inflammatory and host defense  
22 response of human neutrophils. *Eur J Immunol* **40**, 1118-1126, doi:10.1002/eji.200939275 (2010).

23 156 Hiemstra, P. S. Epithelial antimicrobial peptides and proteins: their role in host defence and  
24 inflammation. *Paediatr Respir Rev* **2**, 306-310, doi:10.1053/prrv.2001.0165 (2001).

25 157 Yang, D., Biragyn, A., Kwak, L. W. & Oppenheim, J. J. Mammalian defensins in immunity: more than just  
26 microbicidal. *Trends Immunol* **23**, 291-296 (2002).

27 158 Biragyn, A. *et al.* Toll-like receptor 4-dependent activation of dendritic cells by beta-defensin 2. *Science*  
28 **298**, 1025-1029, doi:10.1126/science.1075565 (2002).

29 159 Davidson, D. J. *et al.* The cationic antimicrobial peptide LL-37 modulates dendritic cell differentiation  
30 and dendritic cell-induced T cell polarization. *J Immunol* **172**, 1146-1156 (2004).

31 160 Tewary, P. *et al.* beta-Defensin 2 and 3 promote the uptake of self or CpG DNA, enhance IFN-alpha  
32 production by human plasmacytoid dendritic cells, and promote inflammation. *J Immunol* **191**, 865-  
33 874, doi:10.4049/jimmunol.1201648 (2013).

34 161 Funderburg, N. *et al.* Human -defensin-3 activates professional antigen-presenting cells via Toll-like  
35 receptors 1 and 2. *Proc Natl Acad Sci U S A* **104**, 18631-18635, doi:0702130104 [pii]  
36 10.1073/pnas.0702130104 (2007).

37 162 Bandholtz, L. *et al.* Antimicrobial peptide LL-37 internalized by immature human dendritic cells alters  
38 their phenotype. *Scand J Immunol* **63**, 410-419 (2006).

39 **163** Findlay, E. G. *et al.* Exposure to the anti-microbial peptide LL-37 produces dendritic cells optimised for  
40 immunotherapy *Oncol Immunology In Press* (2019). **This recent study shows that CHDP can enhance**  
41 **cytotoxic T cells responses to promote tumour clearance in vivo, via effects in dendritic cell**  
42 **differentiation and function, highlighting adjuvant and immunotherapy potential for cancers.**

43 164 Lande, R. *et al.* Plasmacytoid dendritic cells sense self-DNA coupled with antimicrobial peptide. *Nature*  
44 **449**, 564-569, doi:10.1038/nature06116 (2007).

45 165 Ganguly, D. *et al.* Self-RNA-antimicrobial peptide complexes activate human dendritic cells through  
46 TLR7 and TLR8. *J Exp Med* **206**, 1983-1994, doi:10.1084/jem.20090480 (2009).

47 166 Kim, S. H., Kim, Y. N. & Jang, Y. S. Cutting Edge: LL-37-Mediated Formyl Peptide Receptor-2 Signaling in  
48 Follicular Dendritic Cells Contributes to B Cell Activation in Peyer's Patch Germinal Centers. *J Immunol*  
49 **198**, 629-633, doi:10.4049/jimmunol.1600886 (2017).

1 167 Severino, P. *et al.* Cathelicidin-deficient mice exhibit increased survival and upregulation of key  
2 inflammatory response genes following cecal ligation and puncture. *J Mol Med (Berl)* **95**, 995-1003,  
3 doi:10.1007/s00109-017-1555-z (2017).

4 168 Deng, Y. Y., Shamoon, M., He, Y., Bhatia, M. & Sun, J. Cathelicidin-related antimicrobial peptide  
5 modulates the severity of acute pancreatitis in mice. *Mol Med Rep* **13**, 3881-3885,  
6 doi:10.3892/mmr.2016.5008 (2016).

7 169 Wehkamp, J., Schmid, M., Fellermann, K. & Stange, E. F. Defensin deficiency, intestinal microbes, and  
8 the clinical phenotypes of Crohn's disease. *J Leukoc Biol* **77**, 460-465, doi:jlb.0904543 [pii]  
9 10.1189/jlb.0904543 (2005).

10 170 Cirioni, O. *et al.* LL-37 protects rats against lethal sepsis caused by gram-negative bacteria. *Antimicrob*  
11 *Agents Chemother* **50**, 1672-1679 (2006).

12 171 Fukumoto, K. *et al.* Effect of antibacterial cathelicidin peptide CAP18/LL-37 on sepsis in neonatal rats.  
13 *Pediatr Surg Int* **21**, 20-24 (2005).

14 172 Giacometti, A. *et al.* The antimicrobial peptide BMAP-28 reduces lethality in mouse models of  
15 staphylococcal sepsis. *Crit Care Med* **32**, 2485-2490 (2004).

16 **173** Coorens, M. *et al.* Killing of *P. aeruginosa* by chicken cathelicidin-2 is immunogenically silent,  
17 preventing lung inflammation *in vivo*. *Infect Immun*, doi:10.1128/IAI.00546-17 (2017). **This study**  
18 **demonstrates that chicken cathelicidin CATH-2 kills *P.aeruginosa* in an immunogenically silent**  
19 **manner and controls lung inflammation *in vivo*, making a case for the development of CHDP-based**  
20 **anti-infectives with both antimicrobial and immunomodulatory functions.**

21 174 Nagaoka, I. *et al.* Cathelicidin family of antibacterial peptides CAP18 and CAP11 inhibit the expression  
22 of TNF-alpha by blocking the binding of LPS to CD14(+) cells. *J Immunol* **167**, 3329-3338 (2001).

23 175 Rosenfeld, Y., Papo, N. & Shai, Y. Endotoxin (lipopolysaccharide) neutralization by innate immunity  
24 host-defense peptides. Peptide properties and plausible modes of action. *J Biol Chem* **281**, 1636-1643  
25 (2006).

26 **176** Molhoek, E. M. *et al.* Structure-function relationship of the human antimicrobial peptide LL-37 and LL-  
27 **37 fragments in the modulation of TLR responses. *Biol Chem* **390**, 295-303, doi:10.1515/BC.2009.037**  
28 **10.1515/BC.2009.037 [pii] (2009). The first study to interrogate the sequence-to-function relationship in the**  
29 **immunomodulatory functions of CHDP.**

30 177 Choi, K. Y., Napper, S. & Mookherjee, N. Human cathelicidin LL-37 and its derivative IG-19 regulate  
31 interleukin-32-induced inflammation. *Immunology* **143**, 68-80, doi:10.1111/imm.12291 (2014).

32 178 Zhang, Z., Cherryholmes, G. & Shively, J. E. Neutrophil secondary necrosis is induced by LL-37 derived  
33 from cathelicidin. *J Leukoc Biol* **84**, 780-788, doi:10.1189/jlb.0208086 (2008).

34 179 Merkle, M. *et al.* LL37 inhibits the inflammatory endothelial response induced by viral or endogenous  
35 DNA. *J Autoimmun* **65**, 19-29, doi:10.1016/j.jaut.2015.07.015 (2015).

36 180 van Dijk, A. *et al.* Immunomodulatory and Anti-Inflammatory Activities of Chicken Cathelicidin-2  
37 Derived Peptides. *PLoS One* **11**, e0147919, doi:10.1371/journal.pone.0147919 (2016).

38 181 Mookherjee, N. *et al.* Bovine and human cathelicidin cationic host defense peptides similarly suppress  
39 transcriptional responses to bacterial lipopolysaccharide. *J Leukoc Biol* **80**, 1563-1574, doi:jlb.0106048  
40 [pii]  
41 10.1189/jlb.0106048 (2006).

42 182 Coorens, M., van Dijk, A., Bikker, F., Veldhuizen, E. J. & Haagsman, H. P. Importance of Endosomal  
43 Cathelicidin Degradation To Enhance DNA-Induced Chicken Macrophage Activation. *J Immunol* **195**,  
44 3970-3977, doi:10.4049/jimmunol.1501242 (2015).

45 183 Coorens, M. *et al.* Cathelicidins Inhibit *Escherichia coli*-Induced TLR2 and TLR4 Activation in a Viability-  
46 Dependent Manner. *J Immunol* **199**, 1418-1428, doi:10.4049/jimmunol.1602164 (2017).

1 184 McHugh, B. J. *et al.* Cathelicidin is a “fire alarm”, generating protective NLRP3-dependent airway  
2 epithelial cell inflammatory responses during infection with *Pseudomonas aeruginosa*. *PLoS Pathogens*  
3 **In Press** (2019).

4 185 Semple, F. *et al.* Human beta-defensin 3 affects the activity of pro-inflammatory pathways associated  
5 with MyD88 and TRIF. *Eur J Immunol*, doi:10.1002/eji.201141648 (2011).

6 186 Semple, F. *et al.* Human beta-defensin 3 has immunosuppressive activity in vitro and in vivo. *Eur J*  
7 *Immunol* **40**, 1073-1078, doi:10.1002/eji.200940041 (2010).

8 187 McGlasson, S. L. *et al.* Human beta-defensin 3 increases the TLR9-dependent response to bacterial  
9 DNA. *Eur J Immunol* **47**, 658-664, doi:10.1002/eji.201646799 (2017).

10 188 Semple, F. *et al.* Human beta-Defensin 3 [corrected] Exacerbates MDA5 but Suppresses TLR3  
11 Responses to the Viral Molecular Pattern Mimic Polyinosinic:Polycytidylic Acid. *PLoS Genet* **11**,  
12 e1005673, doi:10.1371/journal.pgen.1005673 (2015).

13 189 Funderburg, N. T. *et al.* The Toll-like receptor 1/2 agonists Pam(3) CSK(4) and human beta-defensin-3  
14 differentially induce interleukin-10 and nuclear factor-kappaB signalling patterns in human monocytes.  
15 *Immunology* **134**, 151-160, doi:10.1111/j.1365-2567.2011.03475.x (2011).

16 190 Lande, R. *et al.* Neutrophils activate plasmacytoid dendritic cells by releasing self-DNA-peptide  
17 complexes in systemic lupus erythematosus. *Sci Transl Med* **3**, 73ra19, doi:3/73/73ra19 [pii]  
18 10.1126/scitranslmed.3001180 (2011).

19 191 Filewod, N. C., Pistolic, J. & Hancock, R. E. Low concentrations of LL-37 alter IL-8 production by  
20 keratinocytes and bronchial epithelial cells in response to proinflammatory stimuli. *FEMS Immunol Med*  
21 *Microbiol* **56**, 233-240, doi:FIM571 [pii]  
22 10.1111/j.1574-695X.2009.00571.x (2009).

23 192 Lai, Y. *et al.* LL37 and cationic peptides enhance TLR3 signaling by viral double-stranded RNAs. *PLoS One*  
24 **6**, e26632, doi:10.1371/journal.pone.0026632 (2011).

25 193 Elssner, A., Duncan, M., Gavrillin, M. & Wewers, M. D. A novel P2X7 receptor activator, the human  
26 cathelicidin-derived peptide LL37, induces IL-1 beta processing and release. *J Immunol* **172**, 4987-4994  
27 (2004).

28 194 Li, N. *et al.* Alarmin function of cathelicidin antimicrobial peptide LL37 through IL-36gamma induction  
29 in human epidermal keratinocytes. *J Immunol* **193**, 5140-5148, doi:10.4049/jimmunol.1302574 (2014).

30 195 Yu, J. *et al.* Host defense peptide LL-37, in synergy with inflammatory mediator IL-1beta, augments  
31 immune responses by multiple pathways. *J Immunol* **179**, 7684-7691, doi:179/11/7684 [pii] (2007).

32 196 Lau, Y. E. *et al.* Interaction and cellular localization of the human host defense peptide LL-37 with lung  
33 epithelial cells. *Infect Immun* **73**, 583-591, doi:10.1128/IAI.73.1.583-591.2005 (2005).

34 197 Sandgren, S. *et al.* The human antimicrobial peptide LL-37 transfers extracellular DNA plasmid to the  
35 nuclear compartment of mammalian cells via lipid rafts and proteoglycan-dependent endocytosis. *J*  
36 *Biol Chem* **279**, 17951-17956, doi:10.1074/jbc.M311440200 (2004).

37 198 Lau, Y. E., Bowdish, D. M., Cosseau, C., Hancock, R. E. & Davidson, D. J. Apoptosis of airway epithelial  
38 cells: human serum sensitive induction by the cathelicidin LL-37. *Am J Respir Cell Mol Biol* **34**, 399-409  
39 (2006).

40 199 Barlow, P. G. *et al.* The human cathelicidin LL-37 preferentially promotes apoptosis of infected airway  
41 epithelium. *Am J Respir Cell Mol Biol* **43**, 692-702, doi:2009-0250OC [pii]  
42 10.1165/rcmb.2009-0250OC (2010).

43 200 Barlow, P. G. *et al.* The human cationic host defense peptide LL-37 mediates contrasting effects on  
44 apoptotic pathways in different primary cells of the innate immune system. *J Leukoc Biol* **80**, 509-520  
45 (2006).

1 201 Mader, J. S., Mookherjee, N., Hancock, R. E. & Bleackley, R. C. The human host defense peptide LL-37  
2 induces apoptosis in a calpain- and apoptosis-inducing factor-dependent manner involving Bax activity.  
3 *Mol Cancer Res* **7**, 689-702, doi:1541-7786.MCR-08-0274 [pii]  
4 10.1158/1541-7786.MCR-08-0274 (2009).

5 202 Bowdish, D. M., Davidson, D. J., Speert, D. P. & Hancock, R. E. The human cationic peptide LL-37  
6 induces activation of the extracellular signal-regulated kinase and p38 kinase pathways in primary  
7 human monocytes. *J Immunol* **172**, 3758-3765 (2004).

8 203 Savill, J., Dransfield, I., Gregory, C. & Haslett, C. A blast from the past: clearance of apoptotic cells  
9 regulates immune responses. *Nat Rev Immunol* **2**, 965-975, doi:10.1038/nri957 (2002).

10 204 Li, H. N. *et al.* Secondary necrosis of apoptotic neutrophils induced by the human cathelicidin LL-37 is  
11 not proinflammatory to phagocytosing macrophages. *J Leukoc Biol* **86**, 891-902, doi:jlb.0209050 [pii]  
12 10.1189/jlb.0209050 (2009).

13 205 Miles, K. *et al.* Dying and necrotic neutrophils are anti-inflammatory secondary to the release of alpha-  
14 defensins. *J Immunol* **183**, 2122-2132, doi:10.4049/jimmunol.0804187 (2009).

15 206 Nizet, V. *et al.* Innate antimicrobial peptide protects the skin from invasive bacterial infection. *Nature*  
16 **414**, 454-457 (2001).

17 207 Yu, F. S. *et al.* Flagellin stimulates protective lung mucosal immunity: role of cathelicidin-related  
18 antimicrobial peptide. *J Immunol* **185**, 1142-1149, doi:10.4049/jimmunol.1000509 (2010).

19 208 Huang, L. C., Reins, R. Y., Gallo, R. L. & McDermott, A. M. Cathelicidin-deficient (Cnlp <sup>-/-</sup>) mice show  
20 increased susceptibility to *Pseudomonas aeruginosa* keratitis. *Invest Ophthalmol Vis Sci* **48**, 4498-4508,  
21 doi:10.1167/iovs.07-0274 (2007).

22 209 Iimura, M. *et al.* Cathelicidin mediates innate intestinal defense against colonization with epithelial  
23 adherent bacterial pathogens. *J Immunol* **174**, 4901-4907 (2005).

24 210 Chromek, M. *et al.* The antimicrobial peptide cathelicidin protects the urinary tract against invasive  
25 bacterial infection. *Nat Med* **12**, 636-641, doi:nm1407 [pii]  
26 10.1038/nm1407 (2006).

27 211 Kovach, M. A. *et al.* Cathelicidin-related antimicrobial peptide is required for effective lung mucosal  
28 immunity in Gram-negative bacterial pneumonia. *J Immunol* **189**, 304-311,  
29 doi:10.4049/jimmunol.1103196 (2012).

30 212 Pachon-Ibanez, M. E., Smani, Y., Pachon, J. & Sanchez-Cespedes, J. Perspectives for clinical use of  
31 engineered human host defense antimicrobial peptides. *FEMS Microbiol Rev* **41**, 323-342,  
32 doi:10.1093/femsre/fux012 (2017).

33 213 Woodburn, K. W., Jaynes, J. M. & Clemens, L. E. Evaluation of the Antimicrobial Peptide, RP557, for the  
34 Broad-Spectrum Treatment of Wound Pathogens and Biofilm. *Front Microbiol* **10**, 1688,  
35 doi:10.3389/fmicb.2019.01688 (2019).

36 214 Tenland, E. *et al.* A novel derivative of the fungal antimicrobial peptide plectasin is active against  
37 *Mycobacterium tuberculosis*. *Tuberculosis (Edinb)* **113**, 231-238, doi:10.1016/j.tube.2018.10.008  
38 (2018).

39 215 Obuobi, S. *et al.* Facile and efficient encapsulation of antimicrobial peptides via crosslinked DNA  
40 nanostructures and their application in wound therapy. *J Control Release* **313**, 120-130,  
41 doi:10.1016/j.jconrel.2019.10.013 (2019).

42 216 Riool, M., de Breij, A., Drijfhout, J. W., Nibbering, P. H. & Zaat, S. A. J. Antimicrobial Peptides in  
43 Biomedical Device Manufacturing. *Front Chem* **5**, 63, doi:10.3389/fchem.2017.00063 (2017).

44 217 de Breij, A. *et al.* The antimicrobial peptide SAAP-148 combats drug-resistant bacteria and biofilms. *Sci*  
45 *Transl Med* **10**, doi:10.1126/scitranslmed.aan4044 (2018).

46 218 Jiang, Y. *et al.* Antiviral activity of recombinant mouse beta-defensin 3 against influenza A virus in vitro  
47 and in vivo. *Antivir Chem Chemother* **22**, 255-262, doi:10.3851/IMP2077 (2012).



1 219 Gong, T. *et al.* Recombinant mouse beta-defensin 2 inhibits infection by influenza A virus by blocking its  
2 entry. *Arch Virol* **155**, 491-498, doi:10.1007/s00705-010-0608-1 (2010).

3 220 Li, W. *et al.* Construction of eukaryotic expression vector with mBD1-mBD3 fusion genes and exploring  
4 its activity against influenza A virus. *Viruses* **6**, 1237-1252, doi:10.3390/v6031237 (2014).

5 221 Wohlford-Lenane, C. L. *et al.* Rhesus theta-defensin prevents death in a mouse model of severe acute  
6 respiratory syndrome coronavirus pulmonary disease. *J Virol* **83**, 11385-11390, doi:10.1128/JVI.01363-  
7 09 (2009).

8 222 Greber, K. E. & Dawgul, M. Antimicrobial Peptides Under Clinical Trials. *Curr Top Med Chem* **17**, 620-  
9 628, doi:10.2174/1568026616666160713143331 (2017).

10 223 Wiig, M. E. *et al.* PXL01 in sodium hyaluronate for improvement of hand recovery after flexor tendon  
11 repair surgery: randomized controlled trial. *PLoS One* **9**, e110735, doi:10.1371/journal.pone.0110735  
12 (2014).

13 **224** Gronberg, A., Mahlapuu, M., Stahle, M., Whately-Smith, C. & Rollman, O. Treatment with LL-37 is safe  
14 and effective in enhancing healing of hard-to-heal venous leg ulcers: a randomized, placebo-controlled  
15 clinical trial. *Wound Repair Regen* **22**, 613-621, doi:10.1111/wrr.12211 (2014). **The first randomized  
16 clinical trial using a CHDP shown effective in hard-to-heal venous leg ulcers.**

17 225 Martineau, A. R. *et al.* Vitamin D3 supplementation in patients with chronic obstructive pulmonary  
18 disease (ViDiCO): a multicentre, double-blind, randomised controlled trial. *Lancet Respir Med* **3**, 120-  
19 130, doi:10.1016/S2213-2600(14)70255-3 (2015).

20 226 Mily, A. *et al.* Oral intake of phenylbutyrate with or without vitamin D3 upregulates the cathelicidin LL-  
21 37 in human macrophages: a dose finding study for treatment of tuberculosis. *BMC Pulm Med* **13**, 23,  
22 doi:10.1186/1471-2466-13-23 (2013).

23 227 Lipsky, B. A., Holroyd, K. J. & Zasloff, M. Topical versus systemic antimicrobial therapy for treating  
24 mildly infected diabetic foot ulcers: a randomized, controlled, double-blinded, multicenter trial of  
25 pexiganan cream. *Clin Infect Dis* **47**, 1537-1545, doi:10.1086/593185 (2008).

26 228 Bals, R., Weiner, D. J., Mosconi, A. D., Meegalla, R. L. & Wilson, J. M. Augmentation of innate host  
27 defense by expression of a cathelicidin antimicrobial peptide. *Infect Immun* **67**, 6084-6089 (1999).

28 **229** Cuperus, T., van Dijk, A., Matthijs, M. G., Veldhuizen, E. J. & Haagsman, H. P. Protective effect of in ovo  
29 treatment with the chicken cathelicidin analog D-CATH-2 against avian pathogenic E. coli. *Sci Rep* **6**,  
30 26622, doi:10.1038/srep26622 (2016). **The first study to demonstrate that prophylactic cathelicidin  
31 treatment of chick embryos in ovo partially protects chickens from E.coli infection 10 days later, thus  
32 establishing the use in veterinary medicine.**

33 230 North, J. R. *et al.* A novel approach for emerging and antibiotic resistant infections: Innate defense  
34 regulators as an agnostic therapy. *J Biotechnol* **226**, 24-34, doi:10.1016/j.jbiotec.2016.03.032 (2016).

35 231 Niyonsaba, F. *et al.* The innate defense regulator peptides IDR-HH2, IDR-1002, and IDR-1018 modulate  
36 human neutrophil functions. *J Leukoc Biol*, doi:10.1189/jlb.1012497 [pii]  
37 10.1189/jlb.1012497 (2013).

38 232 Tai, E. K. *et al.* Intrarectal administration of mCRAMP-encoding plasmid reverses exacerbated colitis in  
39 Cnlp(-/-) mice. *Gene Ther* **20**, 187-193, doi:10.1038/gt.2012.22 (2013).

40 233 Tai, E. K. *et al.* A new role for cathelicidin in ulcerative colitis in mice. *Exp Biol Med (Maywood)* **232**,  
41 799-808, doi:10.1177/0740779907301111 [pii] (2007).

42 234 Wong, C. C. *et al.* Protective effects of cathelicidin-encoding Lactococcus lactis in murine ulcerative  
43 colitis. *J Gastroenterol Hepatol* **27**, 1205-1212, doi:10.1111/j.1440-1746.2012.07158.x (2012).

44 235 Hing, T. C. *et al.* The antimicrobial peptide cathelicidin modulates Clostridium difficile-associated colitis  
45 and toxin A-mediated enteritis in mice. *Gut* **62**, 1295-1305, doi:10.1136/gutjnl-2012-302180 (2013).

46 236 Otvos, L. Host Defense Peptides and Cancer; Perspectives on Research Design and Outcomes. *Protein  
47 Pept Lett* **24**, 879-886, doi:10.2174/0929866524666170202153501 (2017).

48 237 Wu, W. K. *et al.* Emerging roles of the host defense peptide LL-37 in human cancer and its potential  
49 therapeutic applications. *Int J Cancer* **127**, 1741-1747, doi:10.1002/ijc.25489 (2010).

1 238 Roudi, R., Syn, N. L. & Roudbary, M. Antimicrobial Peptides As Biologic and Immunotherapeutic Agents  
2 against Cancer: A Comprehensive Overview. *Front Immunol* **8**, 1320, doi:10.3389/fimmu.2017.01320  
3 (2017).

4 239 Schulze, K., Ebensen, T., Babiuk, L. A., Gerds, V. & Guzman, C. A. Intranasal vaccination with an  
5 adjuvanted polyphosphazenes nanoparticle-based vaccine formulation stimulates protective immune  
6 responses in mice. *Nanomedicine* **13**, 2169-2178, doi:10.1016/j.nano.2017.05.012 (2017).

7 240 Garlapati, S. *et al.* Immunization with PCEP microparticles containing pertussis toxoid, CpG ODN and a  
8 synthetic innate defense regulator peptide induces protective immunity against pertussis. *Vaccine* **29**,  
9 6540-6548, doi:10.1016/j.vaccine.2011.07.009 (2011).

10 241 Yu, C. H. *et al.* Synthetic innate defense regulator peptide combination using CpG ODN as a novel  
11 adjuvant induces longlasting and balanced immune responses. *Mol Med Rep* **13**, 915-924,  
12 doi:10.3892/mmr.2015.4581 (2016).

13 242 Sorensen, O. E. *et al.* Wound healing and expression of antimicrobial peptides/polypeptides in human  
14 keratinocytes, a consequence of common growth factors. *J Immunol* **170**, 5583-5589 (2003).

15 243 Baroni, A. *et al.* Antimicrobial human beta-defensin-2 stimulates migration, proliferation and tube  
16 formation of human umbilical vein endothelial cells. *Peptides* **30**, 267-272,  
17 doi:10.1016/j.peptides.2008.11.001 (2009).

18 244 Carretero, M. *et al.* In vitro and in vivo wound healing-promoting activities of human cathelicidin LL-37.  
19 *J Invest Dermatol* **128**, 223-236, doi:5701043 [pii]  
20 10.1038/sj.jid.5701043 (2008).

21 245 Koczulla, R. *et al.* An angiogenic role for the human peptide antibiotic LL-37/hCAP-18. *J Clin Invest* **111**,  
22 1665-1672 (2003).

23 246 Arampatzoglou, A. *et al.* Clarithromycin Enhances the Antibacterial Activity and Wound Healing  
24 Capacity in Type 2 Diabetes Mellitus by Increasing LL-37 Load on Neutrophil Extracellular Traps. *Front*  
25 *Immunol* **9**, 2064, doi:10.3389/fimmu.2018.02064 (2018).

26 247 Wang, B. *et al.* IL-1 $\beta$ -Induced Protection of Keratinocytes against Staphylococcus aureus-Secreted  
27 Proteases Is Mediated by Human beta-Defensin 2. *J Invest Dermatol* **137**, 95-105,  
28 doi:10.1016/j.jid.2016.08.025 (2017).

29 248 Heilborn, J. D. *et al.* The cathelicidin anti-microbial peptide LL-37 is involved in re-epithelialization of  
30 human skin wounds and is lacking in chronic ulcer epithelium. *J Invest Dermatol* **120**, 379-389 (2003).

31 249 Dutta, D. *et al.* Development of Silicone Hydrogel Antimicrobial Contact Lenses with Mel4 Peptide  
32 Coating. *Optom Vis Sci* **95**, 937-946, doi:10.1097/OPX.0000000000001282 (2018).

33 250 van Dijk, A., Hedegaard, C. J., Haagsman, H. P. & Heegaard, P. M. H. The potential for immunoglobulins  
34 and host defense peptides (HDPs) to reduce the use of antibiotics in animal production. *Vet Res* **49**, 68,  
35 doi:10.1186/s13567-018-0558-2 (2018).

36 251 Zimmer, J., Hobkirk, J., Mohamed, F., Browning, M. J. & Stover, C. M. On the Functional Overlap  
37 between Complement and Anti-Microbial Peptides. *Front Immunol* **5**, 689,  
38 doi:10.3389/fimmu.2014.00689 (2014).

39 252 Bucki, R., Sostarecz, A. G., Byfield, F. J., Savage, P. B. & Janmey, P. A. Resistance of the antibacterial  
40 agent ceragenin CSA-13 to inactivation by DNA or F-actin and its activity in cystic fibrosis sputum. *J*  
41 *Antimicrob Chemother* **60**, 535-545, doi:10.1093/jac/dkm218 (2007).

42 253 Pezzulo, A. A. *et al.* Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis  
43 lung. *Nature* **487**, 109-113, doi:10.1038/nature11130 (2012).

44 254 Mallia, P. *et al.* Rhinovirus infection induces degradation of antimicrobial peptides and secondary  
45 bacterial infection in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* **186**, 1117-1124,  
46 doi:10.1164/rccm.201205-0806OC (2012).

47 255 Mercer, D. K. *et al.* Improved Methods for Assessing Therapeutic Potential of Antifungal Agents against  
48 Dermatophytes and Their Application in the Development of NP213, a Novel Onychomycosis Therapy  
49 Candidate. *Antimicrob Agents Chemother* **63**, doi:10.1128/AAC.02117-18 (2019).



- 256 Sakoulas, G., Kumaraswamy, M., Kousha, A. & Nizet, V. Interaction of Antibiotics with Innate Host Defense Factors against Salmonella enterica Serotype Newport. *mSphere* **2**, doi:10.1128/mSphere.00410-17 (2017).
- 257 Le, J. *et al.* Effects of vancomycin versus nafcillin in enhancing killing of methicillin-susceptible Staphylococcus aureus causing bacteremia by human cathelicidin LL-37. *Eur J Clin Microbiol Infect Dis* **35**, 1441-1447, doi:10.1007/s10096-016-2682-0 (2016).
- 258 Kumaraswamy, M. *et al.* Standard susceptibility testing overlooks potent azithromycin activity and cationic peptide synergy against MDR Stenotrophomonas maltophilia. *J Antimicrob Chemother* **71**, 1264-1269, doi:10.1093/jac/dkv487 (2016). **This study demonstrates antibiotic synergy with CHDP thus establishing the potential of novel therapeutic cocktails, showing that antibiotic function is affective by native CHDP expression levels, and making the case for the need of new approaches to MIC testing.**
- 259 Lin, L. *et al.* Azithromycin Synergizes with Cationic Antimicrobial Peptides to Exert Bactericidal and Therapeutic Activity Against Highly Multidrug-Resistant Gram-Negative Bacterial Pathogens. *EBioMedicine* **2**, 690-698, doi:10.1016/j.ebiom.2015.05.021 (2015).
- 260 Peric, M. *et al.* Vitamin D analogs differentially control antimicrobial peptide/"alarmin" expression in psoriasis. *PLoS One* **4**, e6340, doi:10.1371/journal.pone.0006340 (2009).
- 261 Sato-Deguchi, E. *et al.* Topical vitamin D(3) analogues induce thymic stromal lymphopoietin and cathelicidin in psoriatic skin lesions. *Br J Dermatol* **167**, 77-84, doi:10.1111/j.1365-2133.2012.10917.x (2012).
- 262 Peng, L. H. *et al.* Integration of antimicrobial peptides with gold nanoparticles as unique non-viral vectors for gene delivery to mesenchymal stem cells with antibacterial activity. *Biomaterials* **103**, 137-149, doi:10.1016/j.biomaterials.2016.06.057 (2016).
- 263 Casciaro, B. *et al.* Gold-nanoparticles coated with the antimicrobial peptide esculentin-1a(1-21)NH<sub>2</sub> as a reliable strategy for antipseudomonal drugs. *Acta Biomater* **47**, 170-181, doi:10.1016/j.actbio.2016.09.041 (2017).
- 264 Piotrowska, U., Sobczak, M. & Oledzka, E. Current state of a dual behaviour of antimicrobial peptides- Therapeutic agents and promising delivery vectors. *Chem Biol Drug Des* **90**, 1079-1093, doi:10.1111/cbdd.13031 (2017).
- 265 Cao, J. *et al.* Yeast-Based Synthetic Biology Platform for Antimicrobial Peptide Production. *ACS Synth Biol* **7**, 896-902, doi:10.1021/acssynbio.7b00396 (2018).
- 266 Wibowo, D. & Zhao, C. X. Recent achievements and perspectives for large-scale recombinant production of antimicrobial peptides. *Appl Microbiol Biotechnol* **103**, 659-671, doi:10.1007/s00253-018-9524-1 (2019).
- 267 Andersson, D. I., Hughes, D. & Kubicek-Sutherland, J. Z. Mechanisms and consequences of bacterial resistance to antimicrobial peptides. *Drug Resist Updat* **26**, 43-57, doi:10.1016/j.drug.2016.04.002 (2016).
- 268 Joo, H. S., Fu, C. I. & Otto, M. Bacterial strategies of resistance to antimicrobial peptides. *Philos Trans R Soc Lond B Biol Sci* **371**, doi:10.1098/rstb.2015.0292 (2016).
- 269 Sahl, H. G. & Shai, Y. Bacterial resistance to antimicrobial peptides. *Biochim Biophys Acta* **1848**, 3019-3020, doi:10.1016/j.bbamem.2015.08.009 (2015).
- 270 El Shazely, B., Urbanski, A., Johnston, P. R. & Rolff, J. In vivo exposure of insect AMP resistant Staphylococcus aureus to an insect immune system. *Insect Biochem Mol Biol* **110**, 60-68, doi:10.1016/j.ibmb.2019.04.017 (2019). **This study discusses the aspects of bacterial resistance to CHDP; demonstrates that resistance generated to CHDP in the laboratory provides no survival advantage to the bacteria in an insect host environment that is dominated by antimicrobial peptides.**
- 271 Tsomaia, N. Peptide therapeutics: targeting the undruggable space. *Eur J Med Chem* **94**, 459-470, doi:10.1016/j.ejmech.2015.01.014 (2015).
- 272 Kintses, B. *et al.* Phylogenetic barriers to horizontal transfer of antimicrobial peptide resistance genes in the human gut microbiota. *Nat Microbiol* **4**, 447-458, doi:10.1038/s41564-018-0313-5 (2019).

- Swidergall, M. & Ernst, J. F. Interplay between *Candida albicans* and the antimicrobial peptide armory. *Eukaryot Cell* **13**, 950-957, doi:10.1128/EC.00093-14 (2014).
- Nakatsuji, T. *et al.* Antimicrobials from human skin commensal bacteria protect against *Staphylococcus aureus* and are deficient in atopic dermatitis. *Sci Transl Med* **9**, doi:10.1126/scitranslmed.aah4680 (2017).
- Doss, M. *et al.* Interactions of alpha-, beta-, and theta-defensins with influenza A virus and surfactant protein D. *J Immunol* **182**, 7878-7887, doi:10.4049/jimmunol.0804049 (2009).
- Leikina, E. *et al.* Carbohydrate-binding molecules inhibit viral fusion and entry by crosslinking membrane glycoproteins. *Nat Immunol* **6**, 995-1001, doi:10.1038/ni1248 (2005).
- Tripathi, S. *et al.* The human cathelicidin LL-37 inhibits influenza A viruses through a mechanism distinct from that of surfactant protein D or defensins. *J Gen Virol* **94**, 40-49, doi:10.1099/vir.0.045013-0 (2013).
- Openshaw, P. J. M., Chiu, C., Culley, F. J. & Johansson, C. Protective and Harmful Immunity to RSV Infection. *Annu Rev Immunol* **35**, 501-532, doi:10.1146/annurev-immunol-051116-052206 (2017).
- Wiens, M. E. & Smith, J. G. alpha-Defensin HD5 Inhibits Human Papillomavirus 16 Infection via Capsid Stabilization and Redirection to the Lysosome. *MBio* **8**, doi:10.1128/mBio.02304-16 (2017).
- Zhang, L., Lopez, P., He, T., Yu, W. & Ho, D. D. Retraction of an interpretation. *Science* **303**, 467, doi:10.1126/science.303.5657.467b (2004).
- Demirkhanyan, L. H. *et al.* Multifaceted mechanisms of HIV-1 entry inhibition by human alpha-defensin. *J Biol Chem* **287**, 28821-28838, doi:10.1074/jbc.M112.375949 (2012).
- Wu, Z. *et al.* Human neutrophil alpha-defensin 4 inhibits HIV-1 infection in vitro. *FEBS Lett* **579**, 162-166, doi:10.1016/j.febslet.2004.11.062 (2005).
- Chang, T. L., Vargas, J., Jr., DelPortillo, A. & Klotman, M. E. Dual role of alpha-defensin-1 in anti-HIV-1 innate immunity. *J Clin Invest* **115**, 765-773, doi:10.1172/JCI21948 (2005).
- Munk, C. *et al.* The theta-defensin, retrocyclin, inhibits HIV-1 entry. *AIDS Res Hum Retroviruses* **19**, 875-881, doi:10.1089/088922203322493049 (2003).
- Owen, S. M. *et al.* RC-101, a retrocyclin-1 analogue with enhanced activity against primary HIV type 1 isolates. *AIDS Res Hum Retroviruses* **20**, 1157-1165, doi:10.1089/aid.2004.20.1157 (2004).
- Cole, A. L. *et al.* HIV-1 adapts to a retrocyclin with cationic amino acid substitutions that reduce fusion efficiency of gp41. *J Immunol* **176**, 6900-6905 (2006).
- Sun, L. *et al.* Human beta-defensins suppress human immunodeficiency virus infection: potential role in mucosal protection. *J Virol* **79**, 14318-14329, doi:10.1128/JVI.79.22.14318-14329.2005 (2005).
- Lafferty, M. K., Sun, L., Christensen-Quick, A., Lu, W. & Garzino-Demo, A. Human Beta Defensin 2 Selectively Inhibits HIV-1 in Highly Permissive CCR6(+)CD4(+) T Cells. *Viruses* **9**, doi:10.3390/v9050111 (2017).
- Quinones-Mateu, M. E. *et al.* Human epithelial beta-defensins 2 and 3 inhibit HIV-1 replication. *AIDS* **17**, F39-48, doi:10.1097/01.aids.0000096878.73209.4f (2003).
- Bergman, P., Walter-Jallow, L., Broliden, K., Agerberth, B. & Soderlund, J. The antimicrobial peptide LL-37 inhibits HIV-1 replication. *Curr HIV Res* **5**, 410-415 (2007).
- Wong, J. H. *et al.* Effects of cathelicidin and its fragments on three key enzymes of HIV-1. *Peptides* **32**, 1117-1122, doi:10.1016/j.peptides.2011.04.017 (2011).
- He, M. *et al.* Cathelicidin-Derived Antimicrobial Peptides Inhibit Zika Virus Through Direct Inactivation and Interferon Pathway. *Front Immunol* **9**, 722, doi:10.3389/fimmu.2018.00722 (2018).

**Acknowledgements:** NM is supported by the Canadian Institutes of Health Research (CIHR) and Natural Sciences and Engineering Research Council (NSERC) Canada for peptide research. MAA is supported by the Australian Research Council. HPH is supported by NWO-ZonMW and NWO-TTW Perspectief grants. DD is

supported by the British Skin Foundation (026/s/17), Action Medical Research (GN2703) and the Chief Scientist Office (TCS/18/02). Authors also gratefully acknowledge Yolanda Gasper and Mark Bleackley, La Trobe University, for their assistance with tables and figures.

#### **Conflict of interest**

NM is listed as an inventor on patents related to immunomodulatory aspects of host defence peptides and IDR peptides. MA is the Chief Scientific Officer and Director of the start-up company Hexima which has a cationic antimicrobial peptide in clinical trials for treatment of onychomycosis. HH owns stock in start-up company Celestial Therapeutics Inc, and has patents on antimicrobial peptide therapeutics licensed to Zoetis. DD declares no conflict of interest.

#### **Publisher's note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### **TOC**

Naturally-occurring cationic host defence peptides (CHDP), also known as antimicrobial peptides, can control infections by their direct microbicidal properties and by modulating host's immune responses. In addition, certain CHDP can resolve harmful inflammation. Here, Mookherjee et al. assess the emerging potential to therapeutically harness these peptides to treat infectious diseases, chronic inflammatory disorders and wound healing, highlighting current preclinical studies and clinical trials.

#### **Subject categories**

Biological sciences / Drug discovery [URI /631/154]

Health sciences / Anatomy / Haemic and immune systems / Immune system [URI/692/698/1543/1565]

Biological sciences / Immunology / Infectious diseases [URI/631/250/255]

Health sciences / Diseases / Immunological disorders / Inflammatory diseases [URI/692/699/249/2510]

Fig 1




Fig 2




Fig 3




Fig 4

